

## Datasheet: MCA2200P

**BATCH NUMBER 162080**

<b>Description:</b>	MOUSE ANTI C-MYC:HRP
<b>Specificity:</b>	C-MYC
<b>Format:</b>	HRP
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	9E10
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen	▪			
Immunohistology - Paraffin	▪			
ELISA	▪			1/100 - 1/500
Immunoprecipitation			▪	
Western Blotting (1)	▪			1/100 - 1/500

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own systems using appropriate negative/positive controls.

**(1)9E10 recognizes c-myc under non-reducing conditions**

### Target Species

Human

### Species Cross Reactivity

Reacts with: Epitope tag

**N.B.** Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

### Product Form

Purified IgG conjugated to Horseradish Peroxidase (HRP) - liquid

### Preparation

Purified IgG prepared by affinity chromatography on Protein G from tissue culture

	supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.01% Thiomersal
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	Synthetic peptide sequence corresponding to the C-terminal region (residues 408-439) of human c-myc conjugated to keyhole limpet hemocyanin.
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P01106</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">4609</a>    MYC    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	BHLHE39
<b>RRID</b>	AB_324087
<b>Fusion Partners</b>	Spleen cells from immunised BALB/c mice were fused with cells of the SP2/0 myeloma cell line.
<b>Specificity</b>	<p><b>Mouse anti c-myc antibody, clone 9E10</b> detects the p62<sup>c-myc</sup> proto-oncogene protein, which is involved in the regulation of the cell cycle and cell growth. C-myc is primarily located to the cell nucleus, but has also been shown to localized to the cytoplasm in several cell lines (<a href="#">Craig et al. 1993</a>). Overexpression of c-myc has been reported in a wide variety of human cancers (Nesbit <i>et al.</i> 1999).</p> <p>Mouse anti c-myc antibody, clone 9E10 recognizes the sequence EQKLISEEDL and may be used to detect proteins and peptides labelled with molecular tags containing this sequence (Hilpert <i>et al.</i> 2001).</p>
<b>Immunohistology</b>	This product does not require protein digestion pre-treatment of paraffin sections prior to staining This product does not require antigen retrieval using heat treatment prior to staining of paraffin sections.
<b>References</b>	<ol style="list-style-type: none"> <li>1. Evan, G.I. <i>et al.</i> (1985) Isolation of monoclonal antibodies specific for human c-myc proto-oncogene product. <a href="#">Mol Cell Biol. 5 (12): 3610-6.</a></li> <li>2. Spandidos, D.A. <i>et al.</i> (1987) Elevated expression of the myc gene in human benign and malignant breast lesions compared to normal tissue. <a href="#">Anticancer Res. 7 (6): 1299-304.</a></li> <li>3. Borodina, I. <i>et al.</i> (2010) Display of wasp venom allergens on the cell surface of <i>Saccharomyces cerevisiae</i>. <a href="#">Microb Cell Fact. 9: 74.</a></li> <li>4. Groeger, G. <i>et al.</i> (2007) Co-operative Cdc42 and Rho signalling mediates ephrinB-triggered endothelial cell retraction. <a href="#">Biochem J. 404: 23-9.</a></li> </ol>

5. Head, B. *et al.* (2009) Inducible proteolytic inactivation of OPA1 mediated by the OMA1 protease in mammalian cells. [J Cell Biol. 187: 959-66.](#)
6. Hilpert, K. *et al.* (2001) Anti-c-myc antibody 9E10: epitope key positions and variability characterized using peptide spot synthesis on cellulose. [Protein Eng. 14: 803-6.](#)
7. Gohlke, S. *et al.* (2017) *In Vitro* and *In Vivo* Studies on the Structural Organization of Chs3 from *Saccharomyces cerevisiae*. [Int J Mol Sci. 18 \(4\): pii: E702.](#)
8. Gray, P. *et al.* (2010) Identification of a novel human MD-2 splice variant that negatively regulates Lipopolysaccharide-induced TLR4 signaling. [J Immunol. 184: 6359-66.](#)
9. Duriseti, S. *et al.* (2010) Antagonistic anti-urokinase plasminogen activator receptor (uPAR) antibodies significantly inhibit uPAR-mediated cellular signaling and migration. [J Biol Chem. 285: 26878-88.](#)
10. Tan, P.H. *et al.* (2005) Creation of tolerogenic human dendritic cells via intracellular CTLA4: a novel strategy with potential in clinical immunosuppression. [Blood. 106: 2936-43.](#)
11. Wallace, S.W. *et al.* (2010) Cdc42 regulates apical junction formation in human bronchial epithelial cells through PAK4 and Par6B. [Mol Biol Cell. 21: 2996-3006.](#)
12. Rowshanravan, B. *et al.* (2014) RasGAP mediates neuronal survival in *Drosophila* through direct regulation of Rab5-dependent endocytosis. [J Cell Sci. 127: 2849-61.](#)
13. Taylor K *et al.* (2015) Nanocell targeting using engineered bispecific antibodies. [MAbs. 7 \(1\): 53-65.](#)
14. Elders, R.C. *et al.* (2014) Recombinant canine IgE Fc and an IgE Fc-TRAIL fusion protein bind to neoplastic canine mast cells. [Vet Immunol Immunopathol. 159 \(1-2\): 29-40.](#)
15. Sharkey, A.M. *et al.* (2015) Tissue-Specific Education of Decidual NK Cells. [J Immunol. 195 \(7\): 3026-32.](#)
16. McGough, I.J. *et al.* (2014) Identification of molecular heterogeneity in SNX27-retromer-mediated endosome-to-plasma-membrane recycling. [J Cell Sci. 127 \(Pt 22\): 4940-53.](#)
17. Frohnert, C. *et al.* (2014) Importin 7 and Nup358 promote nuclear import of the protein component of human telomerase. [PLoS One. 9 \(2\): e88887.](#)
18. Hage, N. *et al.* (2015) Improved expression and purification of the *Helicobacter pylori* adhesin BabA through the incorporation of a hexa-lysine tag. [Protein Expr Purif. 106: 25-30.](#)
19. Mann, J.K. & Park, S. (2015) Epitope-Specific Binder Design by Yeast Surface Display. [Methods Mol Biol. 1319: 143-54.](#)
20. Paraskevopoulou, V. *et al.* (2019) Introduction of a C-terminal hexa-lysine tag increases thermal stability of the LacDiNac binding adhesin (LabA) exodomain from *Helicobacter pylori*. [Protein Expr Purif. 163: 105446.](#)
21. Lim, H.K. *et al.* (2010) Flow cytometric analysis of genetic FRET detectors containing variable substrate sequences. [Biotechnol Prog. 26 \(6\): 1765-71.](#)
22. Walker, L.M. *et al.* (2009) Efficient recovery of high-affinity antibodies from a single-chain Fab yeast display library. [J Mol Biol. 389 \(2\): 365-75.](#)
23. Matos, J. *et al.* (2013) Cell-cycle kinases coordinate the resolution of recombination intermediates with chromosome segregation. [Cell Rep. 4 \(1\): 76-86.](#)
24. Paraskevopoulou, V. *et al.* (2020) Structural and binding characterization of the LacDiNac-specific adhesin (LabA; HopD) exodomain from *Helicobacter pylori*. [Curr Res Struct Biol. 15 Dec \[Epub ahead of print\].](#)
25. Kalusche, S. *et al.* (2020) Lactobacilli Expressing Broadly Neutralizing Nanobodies

against HIV-1 as Potential Vectors for HIV-1 Prophylaxis? [Vaccines \(Basel\). 8 \(4\) Dec 13 \[Epub ahead of print\].](#)

26. Hollandsworth, H.M. *et al.* (2020) Fluorophore-conjugated *Helicobacter pylori* recombinant membrane protein (HopQ) labels primary colon cancer and metastases in orthotopic mouse models by binding CEA-related cell adhesion molecules. [Transl Oncol. 13 \(12\): 100857.](#)

27. Paraskevopoulou, V. *et al.* (2021) Structural and binding characterization of the LacdiNAc-specific adhesin (LabA; HopD) exodomain from *Helicobacter pylori*. [Curr Res Struct Biol. 3: 19-29.](#)

---

<b>Further Reading</b>	1. Nesbit, C. <i>et al.</i> (1999) MYC oncogenes and human neoplastic disease. <a href="#">Oncogene. 18: 3004-16.</a> 2. Krauß, N. <i>et al.</i> (2008) The structure of the anti-c-myc antibody 9E10 Fab fragment/epitope peptide complex reveals a novel binding mode dominated by the heavy chain hypervariable loops. <a href="#">Proteins. 73: 552-65.</a>
------------------------	--

---

<b>Storage</b>	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.
----------------	---

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

---

<b>Guarantee</b>	12 months from date of despatch
------------------	---------------------------------

---

<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10094 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA2200P">https://www.bio-rad-antibodies.com/SDS/MCA2200P</a> 10094
--------------------------------------	---

---

<b>Regulatory</b>	For research purposes only
-------------------	----------------------------

---

## Related Products

### Recommended Useful Reagents

[AbGUARD® HRP STABILIZER PLUS \(BUF052A\)](#)

[AbGUARD® HRP STABILIZER PLUS \(BUF052B\)](#)

[AbGUARD® HRP STABILIZER PLUS \(BUF052C\)](#)

[TMB CORE \(BUF056A\)](#)

[TMB CORE+ \(BUF062A\)](#)

[TMB SIGNAL+ \(BUF054A\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M396898:220615'

Printed on 25 Mar 2023

---

