Datasheet: MCA2209F BATCH NUMBER 160355

Description:	cription: RAT ANTI MOUSE CD79b:FIT		
Specificity:	CD79b		
Other names:	B29		
Format:	FITC		
Product Type:	Monoclonal Antibody		
Clone:	AT107-2		
lsotype:	lgG1		
Quantity:	0.1 mg		

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-</u> rad-antibodies.com/protocols.						
		Yes No	Not Determined	Suggested Dilution			
	Flow Cytometry (1)			1/50 - 1/100			
	necessarily exclude its a guide only. It is recor system using appropria (1) Membrane perme	use in such procedunt mended that the us ate negative/positive abilization is require	er titrates the antibody	g dilutions are given as for use in their own a. The use of			
Target Species	Mouse						
Species Cross Reactivity	Reacts with: Human, Rat, Pig, Dog, Koala, Tasmanian Devil, Cat Does not react with:Chicken N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.						
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid						
Max Ex/Em	Fluorophore	Excitation Max (nm)					
	Fluorophore		Emission Max (nm)				

Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml
Immunogen	Peptide containing 20 amino acid residues from mouse CD79b conjugated to Keyhole Limpet Hemocyanin (KLH).
External Database Links	UniProt: P15530 Related reagents Entrez Gene: 15985 Cd79b Related reagents
Synonyms	lgb
RRID	AB_324172
Fusion Partners	Spleen cells from immunised rat were fused with cells of the mouse NS-1 myeloma cell line.
Specificity	Rat anti Mouse CD79b antibody, clone AT107-2 recognizes a cytoplasmic region of mouse B-cell antigen receptor complex-associated protein beta chain, also known as B-cell-specific glycoprotein B29, Ig-beta, or Immunoglobulin-associated B29 protein. CD79b is a 228 amino acid, including a 26 signal peptide ~40 kDa type I single pass transmembrane glycoprotein. CD79b is expressed by B lymphocytes and associates with CD79a to form a heterodimer, non-covalently linked to surface immunoglobulin, forming the B-cell receptor (BCR) complex. Rat anti Mouse CD79b antibody, clone AT107-2 also recognizes a homologous region of human CD79b.
Flow Cytometry	Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.
References	 Dornan, D. <i>et al.</i> (2009) Therapeutic potential of an anti-CD79b antibody-drug conjugate, anti-CD79b-vc-MMAE, for the treatment of non-Hodgkin lymphoma. <u>Blood. 114:</u> <u>2721-9.</u> Vendel,A.C. <i>et al</i> (2009) B and T lymphocyte attenuator regulates B cell receptor signaling by targeting Syk and BLNK. <u>J Immunol. 182: 1509-17</u> Poggi, A. <i>et al</i> (2017) Prognostic significance of Ki67 evaluated by flow cytometry in dogs with high-grade B-cell lymphoma. <u>Vet Comp Oncol. 15 (2): 431-40.</u> Kreiss, A. <i>et al.</i> (2009) A histological and immunohistochemical analysis of lymphoid tissues of the Tasmanian devil. <u>Anat Rec (Hoboken). 292: 611-20.</u>

	5. Lau, Q. et al. (2012) Expression and in vitro upregulation of MHCII in koala			
	lymphocytes. Vet Immunol Immunopathol. 147: 35-43.			
	6. Kandasamy S et al. (2014) Prenatal vitamin A deficiency impairs adaptive immune			
	responses to pentavalent rotavirus vaccine (RotaTeq ${ m I\!R}$) in a neonatal gnotobiotic pig			
	model. <u>Vaccine. 32 (7): 816-24.</u>			
	7. Whitney, J.L. et al. (2016) Immunohistochemical Analysis of Leucocyte Subsets in the			
	Sinonasal Mucosa of Cats with Upper Respiratory Tract Aspergillosis. J Comp Pathol. 155			
	<u>(2-3): 130-40.</u>			
	8. McCurdy, P. <i>et al.</i> (2014) Acute lymphoblastic leukemia in a pygmy hippopotamus (<i>Hexaprotodon liberiensis</i>). J Zoo Wildl Med. 45 (4): 906-10.			
	9. Allan, J.N. <i>et al.</i> (2019) CD79b Expression in Richter's Transformation <u>Blood. 134</u>			
	(Supplement 1): 4279.			
	10. Thalheim, L. <i>et al.</i> (2013) Lymphoma immunophenotype of dogs determined by			
	immunohistochemistry, flow cytometry, and polymerase chain reaction for antigen receptor			
	rearrangements. <u>J Vet Intern Med. 27 (6): 1509-16.</u> 11. Ouyang, K. <i>et al.</i> (2015) Evaluation of humoral immune status in porcine epidemic			
	diarrhea virus (PEDV) infected sows under field conditions. <u>Vet Res. 46: 140.</u>			
	12. Bennett, A.L> <i>et al.</i> (2017) Canine acute leukaemia: 50 cases (1989-2014). <u>Vet Comp</u>			
	Oncol. 15 (3): 1101-14.			
	13. Borthwick, C.R. <i>et al.</i> (2019) An Examination of the Development and Localization of			
	Key Immune Cells in Developing Pouch Young of the Red-Tailed Phascogale (<i>Phascogale</i>			
	<i>calura</i>). <u>Anat Rec (Hoboken). 302 (11): 1985-2002.</u>			
	14. Tilly, H. <i>et al.</i> (2019) Polatuzumab vedotin in combination with immunochemotherapy			
	in patients with previously untreated diffuse large B-cell lymphoma: an open-label,			
	non-randomised, phase 1b–2 study <u>The Lancet Oncology. 20 (7): 998-1010.</u>			
	15. Sehn, L.H. et al. (2020) Polatuzumab Vedotin in Relapsed or Refractory Diffuse Large			
	B-Cell Lymphoma. <u>J Clin Oncol. 38 (2): 155-65.</u>			
	16. Herrera, A.F. <i>et al.</i> (2022) Anti-CD79B Antibody-Drug Conjugate DCDS0780A in			
	Patients with B-Cell Non-Hodgkin Lymphoma: Phase 1 Dose-Escalation Study. <u>Clin</u>			
	Cancer Res. Jan 03 [Epub ahead of print].			
Further Reading	1. Piriou-Guzylack, L. (2008) Membrane markers of the immune cells in swine: an update.			
	<u>Vet Res. 39: 54.</u>			
Storago	This product is obimed at employed to make a star of the program and at a disput and stars at			
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at 20°C or respired by the second different temperature at 20°C for			
	-20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for			
	short term use (up to 4 weeks) and store the remaining aliquots at -20°C.			
	Avoid repeated freezing and thawing as this may denature the antibody. Storage in			
	frost-free freezers is not recommended. This product is photosensitive and should be			
	protected from light.			
	P			
Guarantee	12 months from date of despatch			
Health And Safety	Material Safety Datasheet documentation #10041 available at:			
Information	https://www.bio-rad-antibodies.com/SDS/MCA2209F			
	10041			

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-ra	Worldwide d.com	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad	Europe I.com	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com	
To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M404420:220823'						

Printed on 15 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint