

Datasheet: MCA2209T

BATCH NUMBER 151999

Description:	RAT ANTI MOUSE CD79b
Specificity:	CD79b
Other names:	B29
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	AT107-2
Isotype:	IgG1
Quantity:	25 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry (1)	▪			1/500 - 1/1000
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			
Immunoprecipitation	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1) Membrane permeabilization is required for this application. The use of Leucoperm (Product Code [BUF09](#)) is recommended for this purpose.

Target Species

Mouse

Species Cross Reactivity

Reacts with: Human, Rat, Pig, Dog, Koala, Tasmanian Devil, Cat
Does not react with: Chicken

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Purified IgG - liquid

Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	TRIS buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Peptide containing 20 amino acid residues from mouse CD79b conjugated to Keyhole Limpet Hemocyanin (KLH).
External Database Links	<p>UniProt: P15530 Related reagents</p> <p>Entrez Gene: 15985 Cd79b Related reagents</p>
Synonyms	Igb
RRID	AB_1102341
Fusion Partners	Spleen cells from immunised rat were fused with cells of the mouse NS-1 myeloma cell line.
Specificity	Rat anti Mouse CD79b antibody, clone AT107-2 recognizes a cytoplasmic region of mouse B-cell antigen receptor complex-associated protein beta chain, also known as B-cell-specific glycoprotein B29, Ig-beta, or Immunoglobulin-associated B29 protein. CD79b is a 228 amino acid, including a 26 signal peptide ~40 kDa type I single pass transmembrane glycoprotein. CD79b is expressed by B lymphocytes and associates with CD79a to form a heterodimer, non-covalently linked to surface immunoglobulin, forming the B-cell receptor (BCR) complex. Rat anti Mouse CD79b antibody, clone AT107-2 also recognizes a homologous region of human CD79b.
Flow Cytometry	Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.
Western Blotting	This clone is suitable for use in Western blotting applications. We recommend the use of MCA2209 for this purpose.
References	<ol style="list-style-type: none"> Dornan, D. <i>et al.</i> (2009) Therapeutic potential of an anti-CD79b antibody-drug conjugate, anti-CD79b-vc-MMAE, for the treatment of non-Hodgkin lymphoma. Blood. 114: 2721-9. Vendel, A.C. <i>et al.</i> (2009) B and T lymphocyte attenuator regulates B cell receptor signaling by targeting Syk and BLNK. J Immunol. 182: 1509-17 Poggi, A. <i>et al.</i> (2017) Prognostic significance of Ki67 evaluated by flow cytometry in dogs with high-grade B-cell lymphoma. Vet Comp Oncol. 15 (2): 431-40.

4. Kreiss, A. *et al.* (2009) A histological and immunohistochemical analysis of lymphoid tissues of the Tasmanian devil. [Anat Rec \(Hoboken\). 292: 611-20.](#)
5. Lau, Q. *et al.* (2012) Expression and *in vitro* upregulation of MHCII in koala lymphocytes. [Vet Immunol Immunopathol. 147: 35-43.](#)
6. Kandasamy S *et al.* (2014) Prenatal vitamin A deficiency impairs adaptive immune responses to pentavalent rotavirus vaccine (RotaTeq®) in a neonatal gnotobiotic pig model. [Vaccine. 32 \(7\): 816-24.](#)
7. Whitney, J.L. *et al.* (2016) Immunohistochemical Analysis of Leucocyte Subsets in the Sinonasal Mucosa of Cats with Upper Respiratory Tract Aspergillosis. [J Comp Pathol. 155 \(2-3\): 130-40.](#)
8. McCurdy, P. *et al.* (2014) Acute lymphoblastic leukemia in a pygmy hippopotamus (*Hexaprotodon liberiensis*). [J Zoo Wildl Med. 45 \(4\): 906-10.](#)
9. Allan, J.N. *et al.* (2019) CD79b Expression in Richter's Transformation [Blood. 134 \(Supplement 1\): 4279.](#)
10. Thalheim, L. *et al.* (2013) Lymphoma immunophenotype of dogs determined by immunohistochemistry, flow cytometry, and polymerase chain reaction for antigen receptor rearrangements. [J Vet Intern Med. 27 \(6\): 1509-16.](#)
11. Ouyang, K. *et al.* (2015) Evaluation of humoral immune status in porcine epidemic diarrhea virus (PEDV) infected sows under field conditions. [Vet Res. 46: 140.](#)
12. Bennett, A.L. *et al.* (2017) Canine acute leukaemia: 50 cases (1989-2014). [Vet Comp Oncol. 15 \(3\): 1101-14.](#)
13. Borthwick, C.R. *et al.* (2019) An Examination of the Development and Localization of Key Immune Cells in Developing Pouch Young of the Red-Tailed Phascogale (*Phascogale calura*). [Anat Rec \(Hoboken\). 302 \(11\): 1985-2002.](#)
14. Tilly, H. *et al.* (2019) Polatuzumab vedotin in combination with immunochemotherapy in patients with previously untreated diffuse large B-cell lymphoma: an open-label, non-randomised, phase 1b–2 study [The Lancet Oncology. 20 \(7\): 998-1010.](#)
15. Sehn, L.H. *et al.* (2020) Polatuzumab Vedotin in Relapsed or Refractory Diffuse Large B-Cell Lymphoma. [J Clin Oncol. 38 \(2\): 155-65.](#)
16. Herrera, A.F. *et al.* (2022) Anti-CD79B Antibody-Drug Conjugate DCDS0780A in Patients with B-Cell Non-Hodgkin Lymphoma: Phase 1 Dose-Escalation Study. [Clin Cancer Res. Jan 03 \[Epub ahead of print\].](#)

Further Reading 1. Piriou-Guzylack, L. (2008) Membrane markers of the immune cells in swine: an update. [Vet Res. 39: 54.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10057 available at: <https://www.bio-rad-antibodies.com/SDS/MCA2209T>
10057

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16...)	DyLight®800
Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (STAR72...)	HRP
Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR73...)	RPE
Rabbit Anti Rat IgG (STAR21...)	HRP
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight®550 , DyLight®650 , DyLight®800
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin

North & South Tel: +1 800 265 7376**America** Fax: +1 919 878 3751Email: antibody_sales_us@bio-rad.com**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M404423:220823'

Printed on 15 Mar 2024