

Datasheet: MCA2218F

Description:	MOUSE ANTI SHEEP CD25:FITC
Specificity:	CD25
Other names:	IL-2R ALPHA CHAIN
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	9.14
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Sheep						
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid						
Max Ex/Em	<table border="1"> <thead> <tr> <th>Fluorophore</th> <th>Excitation Max (nm)</th> <th>Emission Max (nm)</th> </tr> </thead> <tbody> <tr> <td>FITC</td> <td>490</td> <td>525</td> </tr> </tbody> </table>	Fluorophore	Excitation Max (nm)	Emission Max (nm)	FITC	490	525
Fluorophore	Excitation Max (nm)	Emission Max (nm)					
FITC	490	525					
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant						
Buffer Solution	Phosphate buffered saline						
Preservative	0.09% Sodium Azide						
Stabilisers	1% Bovine Serum Albumin						
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml						

External Database Links

UniProt:

[P26898](#) [Related reagents](#)

Entrez Gene:

[443435](#) IL2RA [Related reagents](#)

RRID	AB_2125624
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS-1 myeloma cell line.
Specificity	<p>Mouse anti sheep CD25, clone 9.14 recognizes the ovine homologue of human CD25, also known as interleukin-2 receptor alpha chain (IL-2Rα), a single pass type I membrane protein expressed by activated ovine T-cells.</p> <p>Ovine Interleukin-2 is a cytokine involved in the proliferation, growth and differentiation of T-cells, B-cells and NK cells and its receptor is composed of 3 subunits, an α chain (CD25), a β chain (CD122) and a γ chain (CD132). A non-covalent association of the α and β subunits is required to form the high affinity receptor for IL-2.</p> <p>Mouse anti sheep CD25, clone 9.14 immunoprecipitates a band of ~47 kDa under reducing conditions, as expected for the mature protein due to high glycosylation, consistent with the observed molecular weight of IL-2Rα in other species including humans (Verhagen et al. 1993).</p> <p>Antibodies to CD4, FoxP3 and CD25 may be used in studies of T regulatory cells (T-regs), a unique subset of T helper cells that function in the control of effector cells vital in preventing autoimmunity (Rocchi et al. 2011).</p> <p>Mouse anti sheep CD25, clone 9.14 is one of a wide range of monoclonal antibodies available from Bio-Rad for ovine research and provides an important tool for the identification of ovine CD25 and may be of use in facilitating further studies of T-regs in this species.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.
References	<ol style="list-style-type: none"> Newland, A. et al. (2004) Ovine dendritic cells transduced with an adenoviral CTLA4eEGFP fusion protein construct induce hyporesponsiveness to allostimulation Immunology 113: 310-7. Gillan, S. et al. (2010) Identification of immune parameters to differentiate disease states among sheep infected with <i>Mycobacterium avium</i> subsp. <i>paratuberculosis</i>. Clin Vaccine Immunol. 17: 108-17. Gillan, S. et al. (2010) Ovine immune parameters following immunisation against <i>Mycobacterium avium</i> ssp. <i>paratuberculosis</i> using a lipid-based live-cell vaccine. Vet Immunol Immunopathol. 137 (1-2): 109-19. Piero, B. et al. (2016) Peripheral Blood and Milk Leukocytes Subsets of Lactating Sarda Ewes It J Anim Sci. 12 (2): e34. Willems, M.G. et al. (2016) Systemic interleukin-2 administration improves lung function and modulates chorioamnionitis-induced pulmonary inflammation in the ovine fetus. Am J Physiol Lung Cell Mol Physiol. 310 (1): L1-7.
Further Reading	1. Rocchi, M.S. et al. (2011) Identification of CD4+CD25 high Foxp3+ T cells in ovine peripheral blood. Vet Immunol Immunopathol. 144 (1-2): 172-7.
Storage	<p>Store at +4°C or at -20°C if preferred.</p> <p>This product should be stored undiluted.</p> <p>Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.</p>

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch
------------------	---------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M366366:200529'

Printed on 11 Aug 2020

© 2020 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)