

## Datasheet: MCA2218GA **BATCH NUMBER 169001**

MOUSE ANTI SHEEP CD25
CD25
IL-2R ALPHA CHAIN
Purified
Monoclonal Antibody
9.14
lgG1
0.1 mg

# **Product Details**

Stabilisers

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.biorad-antibodies.com/protocols.

		0010.			
		Yes	No	Not Determined	Suggested Dilution
	Flow Cytometry				1/10 - 1/50
	Immunohistology - Frozen				
	Immunohistology - Paraffin				
	ELISA				
	Immunoprecipitation				
	Western Blotting				
	Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.				
arget Species	Sheep				
Product Form	Purified IgG - liquid				
Preparation	Purified IgG prepared by supernatant	affinity c	hromatog	raphy on Protein A fror	n tissue culture
Buffer Solution	Phosphate buffered salin	e			
Preservative Stabilisers	0.09% sodium azide (Nal	N <sub>3</sub> )			

Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
External Database Links	UniProt: <u>P26898</u> <u>Related reagents</u> Entrez Gene:
	443435 IL2RA Related reagents
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the mouse NS-1 myeloma cell line.
Specificity	<b>Mouse anti Sheep CD25, clone 9.14</b> recognizes the ovine homologue of human CD25, also known as interleukin-2 receptor alpha chain (IL- $2R\alpha$ ), a single pass type I membrane protein expressed by activated ovine T-cells.
	Ovine Interleukin-2 is a cytokine involved in the proliferation, growth and differentiation of T-cells, B-cells and NK cells and its receptor is composed of 3 subunits, an $\alpha$ chain (CD25), a $\beta$ chain (CD122) and a $\gamma$ chain (CD132). A non-covalent association of the $\alpha$ and $\beta$ subunits is required to form the high affinity receptor for IL-2.
	Mouse anti sheep CD25, clone 9.14 immunoprecipitates a band of ~47 kDa under reducing conditions, as expected for the mature protein due to high glycosylation, consistent with the observed molecular weight of IL-2R $\alpha$ in other species including humans (Verhagen <i>et. al.</i> 1993).
	Antibodies to CD4, FoxP3 and CD25 may be used elucidate properties of T regulatory cells (T-regs), a unique subset of T helper cells that function in the control of effector cells vital in preventing autoimmunity ( <u>Rocchi <i>et. al.</i> 2011</u> ).
	Mouse anti sheep CD25, clone 9.14 is one of a wide range of monoclonal antibodies available from Bio-Rad for ovine research and provides an important tool for the identification of ovine CD25.
Flow Cytometry	Use 10µl of the suggested working dilution to label 1 x $10^6$ cells in 100µl
References	<ol> <li>Newland, A. <i>et al.</i> (2004) Ovine dendritic cells transduced with an adenoviral CTLA4eEGFP fusion protein construct induce hyporesponsiveness to allostimulation <u>Immunology 113: 310-7.</u></li> <li>Gillan, S. <i>et al.</i> (2010) Identification of immune parameters to differentiate disease states among sheep infected with <i>Mycobacterium avium</i> subsp. <i>paratuberculosis</i>. <u>Clin</u> <u>Vaccine Immunol. 17: 108-17.</u></li> <li>Gillan, S. <i>et al.</i> (2010) Ovine immune parameters following immunisation against <i>Mycobacterium avium</i> ssp. <i>paratuberculosis</i> using a lipid-based live-cell vaccine. <u>Vet</u> <u>Immunol Immunopathol. 137 (1-2): 109-19.</u></li> </ol>

	4. Piero, B. <i>et al.</i> (2016) Peripheral Blood and Milk Leukocytes Subsets of Lactating Sarda Ewes It J Anim Sci. 12 (2): e34.
	5. Willems, M.G. <i>et al.</i> (2016) Systemic interleukin-2 administration improves lung function and modulates chorioamnionitis-induced pulmonary inflammation in the ovine fetus. <u>Am J Physiol Lung Cell Mol Physiol. 310 (1): L1-7.</u>
	6. Lebedev, M. <i>et al.</i> (2021) Myeloid-like γδ T cell subset in the immune response to an experimental Rift Valley fever vaccine in sheep <u>Veterinary Immunology and</u> <u>Immunopathology. : 110184.</u>
	7. Curina, G. <i>et al.</i> (2018) Evaluation of immune responses in mice and sheep inoculated with a live attenuated <i>Brucella melitensis.</i> REV1 vaccine produced in bioreactor. <u>Vet</u> <u>Immunol Immunopathol. 198: 44-53.</u>
	8. Wooldridge, A.L. <i>et al.</i> (2019) Maternal allergic asthma during pregnancy alters fetal lung and immune development in sheep: potential mechanisms for programming asthma and allergy. <u>J Physiol. 597 (16): 4251-62.</u>
Further Reading	1. Rocchi, M.S. <i>et al.</i> (2011) Identification of CD4+CD25 high Foxp3+ T cells in ovine peripheral blood. <u>Vet Immunol Immunopathol. 144 (1-2): 172-7.</u>
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.
	Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA2218GA 10040
Regulatory	For research purposes only

## **Related Products**

### **Recommended Secondary Antibodies**

Rabbit Anti Mouse IgG (STAR12...)RPERabbit Anti Mouse IgG (STAR13...)HRPGoat Anti Mouse IgG (H/L) (STAR117...)FITCRabbit Anti Mouse IgG (STAR9...)FITCRecommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

North & South	Tel: +1 800 265 7376	Worldwide
America	Fax: +1 919 878 3751	
	Email: antibody_sales_us@bio-rad.c	com

Tel: +44 (0)1865 852 700 **Europe** Fax: +44 (0)1865 852 739 Email: antibody\_sales\_uk@bio-rad.com Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody\_sales\_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

#### Printed on 29 Aug 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint