

## Datasheet: MCA2219GA

<b>Description:</b>	MOUSE ANTI SHEEP CD44
<b>Specificity:</b>	CD44
<b>Other names:</b>	H-CAM, PGP-1
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	25.32
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.1 mg

### Product Details

#### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen	▪			
Immunohistology - Paraffin (1)			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

**(1) This clone has been reported to react with paraffin-embedded material following alcohol fixation. Please refer to reference Sainte-Marie, G. *et al* for information about this protocol.**

<b>Target Species</b>	Sheep
<b>Species Cross Reactivity</b>	<p>Reacts with: Bovine, Goat, Human</p> <p><b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.</p>

<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% sodium azide (NaN <sub>3</sub> )
<b>Carrier Free</b>	Yes
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	Ovine efferent lymphatic duct lymphocytes
<b>Fusion Partners</b>	Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS-1 myeloma cell line.
<b>Specificity</b>	<p><b>Mouse anti Sheep CD44 antibody, clone 25.32</b> recognises the ovine CD44 cell surface antigen, a ~95 kDa glycoprotein expressed by most leucocytes and a subpopulation of thymocytes. In immunohistochemical staining, this antibody labels several cell types including medullary thymocytes, kupffer cells, kidney tubular epithelial cells and gut smooth muscle cells. The expression of CD44 is upregulated upon cell activation.</p> <p>Mouse anti Sheep CD44 antibody, clone 25.32 is a valuable reagent for the isolation and characterization of ovine mesenchymal stem cells along with CD29 and <a href="#">CD166</a> which are also expressed on this cell type, in contrast to hematopoietic sell markers including <a href="#">CD45</a> which is negative on this mesenchymal stem sell population (<a href="#">Sanjurjo-Rodríguez <i>et al.</i> 2017</a>).</p>
<b>Flow Cytometry</b>	Use 10µl of the suggested working dilution to label 1 x 10 <sup>6</sup> cells in 100µl
<b>References</b>	<ol style="list-style-type: none"> <li>1. Aleksandersen, M. <i>et al.</i> (1990) Distribution of lymphocyte subsets in the large intestinal lymphoid follicles of lambs. <a href="#">Immunology. 70 (3): 391-7.</a></li> <li>2. Perry, K. <i>et al.</i> (2010) Hyaluronan (HA) content, the ratio of HA fragments and the expression of CD44 in the ovine cervix vary with the stage of the oestrous cycle. <a href="#">Reproduction. 140:133-41.</a></li> <li>3. Stevenson, L.M. <i>et al.</i> (2001) Expression of cell surface adhesion molecules by peripheral blood eosinophils during <i>Trichostrongylus colubriformis</i> infection in sheep. <a href="#">Immunol Cell Biol. 79 (3): 240-4.</a></li> <li>4. Witherden, D.A. <i>et al.</i> (1995) Antigen-independent maturation of CD2, CD11a/CD18, CD44, and CD58 expression on thymic emigrants in fetal and postnatal sheep. <a href="#">Dev Immunol. 4:199-209</a></li> <li>5. Sanjurjo-Rodríguez, C. <i>et al.</i> (2017) Ovine Mesenchymal Stromal Cells: Morphologic, Phenotypic and Functional Characterization for Osteochondral Tissue Engineering. <a href="#">PLoS One. 12 (1): e0171231.</a></li> <li>6. Wooldridge, A.L. <i>et al.</i> (2019) Maternal allergic asthma during pregnancy alters fetal</li> </ol>

lung and immune development in sheep: potential mechanisms for programming asthma and allergy. [J Physiol. 597 \(16\): 4251-62.](#)

7. López-Fernández, A. *et al.* (2020) Effect of Allogeneic Cell-Based Tissue-Engineered Treatments in a Sheep Osteonecrosis Model. [Tissue Eng Part A. 26 \(17-18\): 993-1004.](#)

8. Zhang, P. *et al.* (2021) Differences in the biological properties of mesenchymal stromal cells from traumatic temporomandibular joint fibrous and bony ankylosis: a comparative study [Animal Cells and Systems. : 1-16.](#)

9. Savy, V. *et al.* (2021) Effect of Embryo Aggregation on *In Vitro* Development of Adipose-Derived Mesenchymal Stem Cell-Derived Bovine Clones. [Cell Reprogram. 23 \(5\): 277-289.](#)

10. Castillo, M.G. *et al.* (2023) Promoting early neovascularization by allotransplanted adipose-derived Muse cells in an ovine model of acute myocardial infarction. [PLoS One. 18 \(1\): e0277442.](#)

---

**Storage** This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

---

**Guarantee** 12 months from date of despatch

---

**Health And Safety Information** Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

---

**Regulatory** For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...) [RPE](#)

Goat Anti Mouse IgG (H/L) (STAR117...) [FITC](#)

Rabbit Anti Mouse IgG (STAR13...) [HRP](#)

Rabbit Anti Mouse IgG (STAR9...) [FITC](#)

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M413351:221121'

Printed on 26 Jan 2023