

# Datasheet: MCA2226F BATCH NUMBER 157535

Description:	MOUSE ANTI SHEEP MHC CLASS II DR MONOMORPHIC:FITC
Specificity:	MHC CLASS II DR MONOMORPHIC
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	37.68
lsotype:	IgG2a
Quantity:	0.1 mg

## **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .					
			No	Not Determined	Suggested Dilution	
	Flow Cytometry	-			1/5 - 1/10	
	Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.					
Target Species	Sheep					
Species Cross Reactivity	Reacts with: Goat, Human, Bovine <b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.					
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid					
Max Ex/Em	Fluorophore FITC	Excitation Max 490	x (nm)	Emission Max (nm) 525		
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant					
Buffer Solution	Phosphate buffered sal	line				

Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml
Immunogen	Splenocytes from ATL mice.
RRID	AB_323966
Specificity	<b>Mouse anti Sheep MHC class II antibody, clone 37.68</b> recognizes a monomorphic epitope on ovine MHC class II DR molecules, constitutively expressed on antigen presenting cells such as dendritic cells, B lymphocytes, monocytes, macrophages, activated T lymphocytes and may be induced on a range of other cell types by interferon gamma.
	The major histocompatibility complex (MHC) is a cluster of genes some of which are important in the immune response to infections. In sheep, this complex is referred to as the ovine leukocyte antigen (OLA) region. There are 2 major types of MHC class IIa molecules encoded by the OLA which are DR and DQ each composed of an alpha and beta chain.
Flow Cytometry	Use 10ul of the suggested working dilution to label $1 \times 10^6$ cells in 100ul.
References	<ol> <li>Puri, N.K. <i>et al.</i> (1985) Sheep lymphocyte antigens (OLA). II. Major histocompatibility complex class II molecules. Immunology. 56 (4): 725-33.</li> <li>Puri, N.K. <i>et al.</i> (1987) Monoclonal antibodies to sheep MHC class I and class II molecules: biochemical characterization of three class I gene products and four distinct subpopulations of class II molecules. <u>Vet Immunol Immunopathol.</u> 15 (1-2): 59-86.</li> <li>Puri, N.K. &amp; Brandon, M.R. (1987) Sheep MHC class II molecules. II. Identification and characterization of four distinct subsets of sheep MHC class II molecules. <u>Immunology.</u> 62 (4): 575-80.</li> <li>Puri, N.K. <i>et al.</i> (1987) Monoclonal antibodies to sheep MHC class II molecules recognize all HLA-D or subsets of HLA-D region products. <u>Hum Immunol.</u> 20 (3): 195-207.</li> <li>Ballingall, K.T. <i>et al.</i> (1995) Analysis of the fine specificities of sheep major histocompatibility complex class II-specific monoclonal antibodies using mouse L-cell transfectants. <u>Anim Genet.</u> 26 (2): 79-84.</li> <li>Wang, Y. <i>et al.</i> (2017) Characterization of a secreted cystatin of the parasitic nematode <i>Haemonchus contortus</i> and its immune-modulatory effect on goat monocytes. <u>Parasit Vectors.</u> 10 (1): 425.</li> <li>Wang, Y. <i>et al.</i> (2020) Characterization of a rhodanese homologue from <i>Haemonchus contortus</i> and its immune-modulatory effects on goat monocytes. <u>Parasit Vectors.</u> 13 (1): 454.</li> <li>Ehsan, M. <i>et al.</i> (2021) <i>Fasciola gigantica.</i> tegumental calcium-binding EF-hand protein 4 exerts immunomodulatory effects on goat monocytes. <u>Parasit Vectors.</u> 14 (1): 276.</li> <li>López-Fernández, A. <i>et al.</i> (2020) Effect of Allogeneic Cell-Based Tissue-Engineered Treatments in a Sheep Osteonecrosis Model. <u>Tissue Eng Part A.</u> 26 (17-18): 993-1004.</li> </ol>

Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.			
	Avoid repeated freezing and thawing as this may denature the frost-free freezers is not recommended. This product is photose protected from light.	, ,		
Guarantee	12 months from date of despatch			
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA2226F 10041			
Regulatory	For research purposes only			

### **Related Products**

#### **Recommended Negative Controls**

MOUSE IgG2a NEGATIVE CONTROL:FITC (MCA929F)

North & South	Tel: +1 800 265 7376 Worldwid	ide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad.com		Email: antibody_sales_uk@bio-rad.com		Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M390544:210921'

#### Printed on 15 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint