

# Datasheet: MCA2228F

Description:	MOUSE ANTI SHEEP MHC CLASS II DQ DR POLYMORPHIC:FITC
Specificity:	MHC CLASS II DQ DR POLYMORPHIC
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	49.1
lsotype:	lgG2a
Quantity:	0.1 mg

### **Product Details**

Applications This product has been reported to work in the following applications. This in						
	derived from testing within our laboratories, peer-reviewed publications or personal					
	communications from the originators. Please refer to references indicated for further					
	information. For generation	al protocol red	commen	dations, please visit <u>w</u>	ww.bio-	
	rad-antibodies.com/protocols.					
		Yes	No	Not Determined	Suggested Dilution	
	Flow Cytometry				1/10	
	Where this product has not been tested for use in a particular technique this does not					
	necessarily exclude its use in such procedures. Suggested working dilutions are given a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.					
Target Species	Sheep					
Species Cross	Reacts with: Goat, Hur	man, Bovine				
Reactivity	Reactivity N.B. Antibody reactivity and working conditions may vary between species. C					
	reactivity is derived fro	m testing with	nin our la	boratories, peer-revie	wed publications or	
	personal communication	ons from the o	originato	rs. Please refer to refe	erences indicated for	
	further information.					
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid					
Max Ex/Em	Fluorophore	Excitation Ma	ax (nm)	Emission Max (nm)		
	FITC	490		525		
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture					
	supernatant					
Buffer Solution	Phosphate buffered sa	line				

0.09% sodium azide (NaN <sub>3</sub> )
1% bovine serum albumin
IgG concentration 0.1 mg/ml
AB_567329
<b>Mouse anti Sheep MHC class II Dq DR antibody, clone 49.1</b> recognizes a polymorphic epitope on ovine MHC class II DQ and DR molecules, constitutively expressed on antigen presenting cells including dendritic cells, B lymphocytes, monocytes, macrophages, activated T lymphocytes and may be induced on a range of other cell types by interferon gamma.
The major histocompatibility complex (MHC) is a cluster of genes some of which are important in the immune response to infection. In sheep, this complex is referred to as the ovine leukocyte antigen (OLA) region. There are 2 major types of MHC class IIa molecules encoded by the OLA which are DR and DQ each composed of an alpha and beta chain.
Mouse anti Sheep MHC class II Dq DR antibody, clone 49.1 recognizes ovine MHC II transfectants DQ - T28.1, DQ - T26.2 and DR - T31.3 but not DR - T8.1 ( <u>Ballingall <i>et al.</i></u> <u>1995</u> ).
Use 10µl of the suggested working dilution to label 1 x $10^6$ cells in 100µl
<ol> <li>Puri, N.K. <i>et al.</i> (1987) Monoclonal antibodies to sheep MHC class I and class II molecules: biochemical characterization of three class I gene products and four distinct subpopulations of class II molecules. <u>Vet Immunol Immunopathol. 15 (1-2): 59-86.</u></li> <li>Puri, N.K. &amp; Brandon, M.R. (1987) Sheep MHC class II molecules. II. Identification and characterization of four distinct subsets of sheep MHC class II molecules. <u>Immunology. 62</u> (4): 575-80.</li> <li>Puri, N.K. <i>et al.</i> (1987) Monoclonal antibodies to sheep MHC class II molecules recognize all HLA-D or subsets of HLA-D region products. <u>Hum Immunol. 20 (3): 195-207.</u></li> <li>Ballingall, K.T. <i>et al.</i> (1995) Analysis of the fine specificities of sheep major histocompatibility complex class II-specific monoclonal antibodies using mouse L-cell transfectants. <u>Anim Genet. 26 (2): 79-84.</u></li> <li>Kallapur, S.G. <i>et al.</i> (2011) Pulmonary and systemic inflammatory responses to intraamniotic IL-1α in fetal sheep. <u>Am J Physiol Lung Cell Mol Physiol. 301 (3): L285-95.</u></li> <li>Gorrell, M.D. <i>et al.</i> (1988) Lymphocyte phenotypes in the intestinal mucosa of sheep infected with <i>Trichostrongylus colubriformis</i>. <u>Clin Exp Immunol. 72 (2): 274-9.</u></li> <li>Stenfeldt, C. <i>et al.</i> (2015) Pathogenesis of Primary Foot-and-Mouth Disease Virus Infection in the Nasopharynx of Vaccinated and Non-Vaccinated Cattle. <u>PLoS One. 10 (11): e0143666.</u></li> <li>Stenfeldt, C. <i>et al.</i> (2016) The Foot-and-Mouth Disease Carrier State Divergence in</li> </ol>

	Cattle. <u>J Virol. 90 (14): 6344-64.</u> 10. Stenfeldt, C. <i>et al.</i> (2019) Virulence beneath the fleece; a ta disease virus pathogenesis in sheep. <u>PLoS One. 14 (12): e022</u>	ale of foot-and-mouth 7061.	
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.		
	Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.		
Guarantee	12 months from date of despatch		
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA2228F">https://www.bio-rad-antibodies.com/SDS/MCA2228F</a> 10041		
Regulatory	For research purposes only		

## Related Products

#### **Recommended Negative Controls**

MOUSE IgG2a NEGATIVE CONTROL:FITC (MCA929F)

North & South	Tel: +1 800 265 7376 Worl	ldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad.com		Email: antibody_sales_uk@bio-rad	.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M413390:221122'

#### Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint