

Datasheet: MCA2238
BATCH NUMBER 153532

Description:	MOUSE ANTI HUMAN CD160
Specificity:	CD160
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	BY55
Isotype:	IgM
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			1/10 - 1/50
Immunohistology - Frozen				
Immunohistology - Paraffin				
ELISA				
Immunoprecipitation	•			
Western Blotting				

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgM - liquid
Preparation	Purified IgM prepared from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃) 1% Bovine Serum Albumin
Approx. Protein Concentrations	IgM concentration 0.1 mg/ml

Immunogen External Database Links	Human NK cell line YT2C2.
	Ha:Dast.
	UniProt: O95971 Related reagents
	Entrez Gene: 11126 CD160 Related reagents
Synonyms	BY55
RRID	AB_2244404
Fusion Partners	Spleen cells from immunized Balb/c mice were fused with cells of the mouse NS-1 myeloma.
Specificity	Mouse anti Human CD160 antibody, clone BY55 recognizes human CD160, a ~27 kDa cell surface glycoprotein that was initially identified by clone BY55. The expression of CD160 is restricted to circulating NK cells, lymphocytes expressing T-cell receptor (TCR) gamma/delta and a small population of TCR alpha/beta positive T cells.
	CD160 is expressed by intestinal intraepithelial lymphocytes.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	 Anumanthan, A. et al. (1998) Cloning of BY55, a novel Ig superfamily member expressed on NK cells, CTL, and intestinal intraepithelial lymphocytes. <u>J Immunol. 161</u> (6): 2780-90. Nikolova, M. et al. (2002) BY55/CD160 acts as a co-receptor in TCR signal transduction of a human circulating cytotoxic effector T lymphocyte subset lacking CD28 expression. <u>Int Immunol. 14 (5): 445-51.</u>
Storage	Store at +4°C or at -20°C if preferred.
	This product should be stored undiluted.
	Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA2238 10041
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgM (STAR138...) Alk. Phos.
Goat Anti Mouse IgG IgA IgM (STAR87...) HRP

Recommended Negative Controls

MOUSE IgM NEGATIVE CONTROL (MCA692)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M366412:200529'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint