

Datasheet: MCA2240F

BATCH NUMBER 166953

Description:	MOUSE ANTI CHICKEN CD41/CD61:FITC
Specificity:	CD41/CD61
Other names:	GPIIb-IIIa
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	11C3
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/2

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Chicken						
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid						
Max Ex/Em	<table border="1"> <thead> <tr> <th>Fluorophore</th> <th>Excitation Max (nm)</th> <th>Emission Max (nm)</th> </tr> </thead> <tbody> <tr> <td>FITC</td> <td>490</td> <td>525</td> </tr> </tbody> </table>	Fluorophore	Excitation Max (nm)	Emission Max (nm)	FITC	490	525
Fluorophore	Excitation Max (nm)	Emission Max (nm)					
FITC	490	525					
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant						
Buffer Solution	Phosphate buffered saline						
Preservative	0.09% sodium azide (NaN ₃)						
Stabilisers	1% bovine serum albumin						
Approx. Protein	IgG concentration 0.1 mg/ml						

Concentrations

Immunogen Peripheral blood leukocytes from white leghorn chicken

RRID AB_2128896

Fusion Partners Spleen cells from immunised BALB/c mice were fused with cells of the X63/A63-653 myeloma cell line

Specificity **Mouse anti Chicken CD41/61, clone 11C3** recognizes chicken integrin CD41/61, also known as GPIIb-IIIa. CD41/61 is a heterodimeric cell surface protein, expressed on chicken thrombocytes and cells of the thrombocyte lineage. Avian thrombocytes are nucleated equivalents of mammalian platelets playing roles in coagulation and phagocytosis.

Mouse anti Chicken CD41/61 clone 11C3 is reported to induce thrombocyte activation ([Lacoste-Eleaume *et al.* 1994](#)).

Flow Cytometry Use 10 μ l of the suggested working dilution to label 10⁶ cells in 100 μ l

References

1. Horiuchi, H. *et al.* (2004) A monoclonal antibody against chicken thrombocytes reacts with the cells of thrombocyte lineage. [J Vet Med Sci. 66 \(3\): 243-50.](#)
 2. Corbel, C. (2002) Expression of alphaVbeta3 integrin in the chick embryo aortic endothelium. [Int J Dev Biol. 46 \(6\): 827-30.](#)
 3. Del Cacho, E. *et al.* (2008) Isolation of chicken follicular dendritic cells. [J Immunol Methods. 334: 59-69.](#)
 4. Del Cacho, E. *et al.* (2009) Avian follicular and interdigitating dendritic cells: isolation and morphologic, phenotypic, and functional analyses. [Vet Immunol Immunopathol. 129: 66-75.](#)
 5. Ody, C. *et al.* (1999) Glycoprotein IIb-IIIa is expressed on avian multilineage hematopoietic progenitor cells. [Blood 93: 2898-906.](#)
 6. Chaussé, A.M. *et al.* (2011) Expression of Toll-like receptor 4 and downstream effectors in selected cecal cell subpopulations of chicks resistant or susceptible to *Salmonella* carrier state. [Infect Immun. 79: 3445-54.](#)
 7. Balic, A. *et al.* (2014) Visualisation of chicken macrophages using transgenic reporter genes: insights into the development of the avian macrophage lineage. [Development 141: 3255-65.](#)
 8. Meijerink, N. *et al.* (2020) Early Life Inoculation With Adult-Derived Microbiota Accelerates Maturation of Intestinal Microbiota and Enhances NK Cell Activation in Broiler Chickens. [Front Vet Sci. 7: 584561.](#)
 9. Meijerink, N. *et al.* (2021) Analysis of chicken intestinal natural killer cells, a major IEL subset during embryonic and early life. [Dev Comp Immunol. 114: 103857.](#)
 10. Meijerink, N. *et al.* (2021) Glucose Oligosaccharide and Long-Chain Glucosaminan Feed Additives Induce Enhanced Activation of Intraepithelial NK Cells and Relative Abundance of Commensal Lactic Acid Bacteria in Broiler Chickens. [Vet Sci. 8 \(6\): 110.](#)
-

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for

short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee	12 months from date of despatch
------------------	---------------------------------

Acknowledgements	This antibody is licensed from INRA-INSERM
-------------------------	--

Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA2240F 10041
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
"M426039:231107"

Printed on 19 Feb 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)