

Datasheet: MCA2243F

BATCH NUMBER 151247

Description:	MOUSE ANTI HUMAN KIR:FITC
Specificity:	KIR
Other names:	KILLER CELL IMMUNOGLOBIN-LIKE RECEPTORS
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	NKVFS1
Isotype:	lgG1
Quantity:	0.1 mg

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .				
		Yes No	Not Determined	Suggested Dilution	
	Flow Cytometry	•		Neat - 1/5	
	Where this antibody has not been tested for use in a particular technique this does no necessarily exclude its use in such procedures. Suggested working dilutions are giver a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.				
Target Species	Human				
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid				
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)		
	FITC	490	525		
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant				
Buffer Solution	Phosphate buffered saline				
Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum Albumin				
Approx. Protein	IgG concentration 0.1 mg/ml				

Concentrations

External Database						
Links	UniProt:					
	<u>P43626</u>	Related reagents				
	<u>P43627</u>	Related reagents				
	<u>P43628</u>	Related reagents				
	<u>Q14954</u>	Related reagents				
	<u>P43631</u>	Related reagents				
	<u>P43632</u>	Related rea	Related reagents			
	Entrez Gene:					
	<u>3802</u>	KIR2DL1	Related reagents			
	<u>3803</u>	KIR2DL2	Related reagents			
	<u>3804</u>	KIR2DL3	Related reagents			
	<u>3806</u>	KIR2DS1	Related reagents			
	<u>3809</u>	KIR2DS4	Related reagents			
	<u>100132285</u>	KIR2DS2	Related reagents			
Synonyms	CD158A, CD158B1, CD158B2, CD158H, CD158I, CD158J, KIRCL23, KKA3, NKAT1, NKAT2, NKAT5, NKAT6, NKAT8					
RRID	AB_324146					
Specificity	Mouse anti Human KIR antibody, clone NKVFS1 recognizes KIR2D members of the killer cell immunoglobulin (Ig)-like receptor (KIR) family, CD158a, CD158b and P50.3. KIR2D family members are cell surface glycoproteins with two Ig domains, which are expressed on natural killer cells and some T cells.					
		Mouse anti Human KIR antibody, clone NKVFS1 recognizes the long and short forms CD158a and CD158b (KIR2DL, KIR2DS1 and KIR2DS2 respectively) and also p50.3 (KIR2DS4).				
	Mouse anti Human KIR antibody, clone NKVFS1 is reported to have functional activity activating NK cell cytotoxicity via KIR2DS and inhibiting via KIR2DL forms					
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.					
References	 Spaggiari, G.M. <i>et al.</i> (2002) Soluble HLA class I molecules induce natural killer cell apoptosis through the engagement of CD8: evidence for a negative regulation exerted by members of the inhibitory receptor superfamily. <u>Blood. 99 (5): 1706-14.</u> Spaggiari, G.M. <i>et al.</i> (2002) Soluble HLA class I induces NK cell apoptosis upon the engagement of killer-activating HLA class I receptors through FasL-Fas interaction. <u>Blood.</u> <u>100 (12): 4098-107.</u> Older Aguilar, A.M. <i>et al.</i> (2010) Coevolution of killer cell Ig-like receptors with HLA-C to become the major variable regulators of human NK cells. <u>J Immunol. 185 (7): 4238-51.</u> Patterson, S. <i>et al.</i> (2008) Human invariant NKT cells display alloreactivity instructed by 					

	Avoid repeated freezing and thawing as this may denature the antibody. Should this
	Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.
	This product should be stored undiluted.
Storage	Store at +4°C or at -20°C if preferred.
	hepatitis B patients. Liver Int. Dec 23 [Epub ahead of print].
	with chronic hepatitis B. <u>Immunol Invest. 38 (5): 434-46.</u> 9. Wijaya, R.S. <i>et al.</i> (2020) Expansion of dysfunctional CD56-CD16+ NK cells in chronic
	8. Wang, Y. et al. (2009) Characteristics of expanded CD4+CD28null T cells in patients
	Cytomegalovirus Stimulates KIR2DS1 Recognition by Natural Killer Cells. <u>Front Immunol.</u> 8: 298.
	7. Van Der Ploeg, K. <i>et al.</i> (2017) Modulation of Human Leukocyte Antigen-C by Human
	epitopes. <u>J Immunol. 182 (6): 3628-37.</u>
	6. Moesta, A.K. <i>et al.</i> (2009) Chimpanzees use more varied receptors and ligands than humans for inhibitory killer cell Ig-like receptor recognition of the MHC-C1 and MHC-C2
	multiplex HLA class I binding assay. <u>J Immunol Methods. 425: 79-87.</u>
	5. Hilton, H.G. et al. (2015) The production of KIR-Fc fusion proteins and their use in a

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL:FITC (MCA928F)

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South	Tel: +1 800 265 7376	Vorldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad.co	om	Email: antibody_sales_uk@bio-rad	d.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M366426:200529'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint