

Datasheet: MCA2243PET

Description:	MOUSE ANTI HUMAN KIR:RPE
Specificity:	KIR
Other names:	KILLER CELL IMMUNOGLOBIN-LIKE RECEPTORS
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	NKVFS1
Isotype:	IgG1
Quantity:	25 TESTS

Product Details

RRID AB_2111691

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species Human

Product Form Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized

Reconstitution Reconstitute in 0.25 ml distilled water

Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE 488nm laser	496	578

Preparation Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

Buffer Solution Phosphate buffered saline

Preservative 0.09% Sodium Azide
Stabilisers 1% Bovine Serum Albumin
 5% Sucrose

External Database Links

UniProt:
[P43626](#) [Related reagents](#)

[P43627](#) [Related reagents](#)
[P43628](#) [Related reagents](#)
[Q14954](#) [Related reagents](#)
[P43631](#) [Related reagents](#)
[P43632](#) [Related reagents](#)

Entrez Gene:

[3802](#) KIR2DL1 [Related reagents](#)
[3803](#) KIR2DL2 [Related reagents](#)
[3804](#) KIR2DL3 [Related reagents](#)
[3806](#) KIR2DS1 [Related reagents](#)
[3809](#) KIR2DS4 [Related reagents](#)
[100132285](#) KIR2DS2 [Related reagents](#)

Synonyms CD158A, CD158B1, CD158B2, CD158H, CD158I, CD158J, KIRCL23, KKA3, NKAT1, NKAT2, NKAT5, NKAT6, NKAT8

Specificity **Mouse anti Human KIR antibody, clone NKVFS1** recognizes KIR2D members of the killer cell immunoglobulin (Ig)-like receptor (KIR) family, CD158a, CD158b and P50.3. KIR2D family members are cell surface glycoproteins with two Ig domains, which are expressed on natural killer cells and some T cells.

Mouse anti Human KIR antibody, clone NKVFS1 recognizes the long and short forms CD158a and CD158b (KIR2DL, KIR2DS1 and KIR2DS2 respectively) and also p50.3 (KIR2DS4).

Mouse anti Human KIR antibody, clone NKVFS1 is reported to have functional activity, activating NK cell cytotoxicity via KIR2DS and inhibiting via KIR2DL forms..

Flow Cytometry Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

References

1. Spaggiari, G.M. *et al.* (2002) Soluble HLA class I molecules induce natural killer cell apoptosis through the engagement of CD8: evidence for a negative regulation exerted by members of the inhibitory receptor superfamily. [Blood. 99 \(5\): 1706-14.](#)
2. Spaggiari, G.M. *et al.* (2002) Soluble HLA class I induces NK cell apoptosis upon the engagement of killer-activating HLA class I receptors through FasL-Fas interaction. [Blood. 100 \(12\): 4098-107.](#)
3. Older Aguilar, A.M. *et al.* (2010) Coevolution of killer cell Ig-like receptors with HLA-C to become the major variable regulators of human NK cells. [J Immunol. 185 \(7\): 4238-51.](#)
4. Patterson, S. *et al.* (2008) Human invariant NKT cells display alloreactivity instructed by invariant TCR-CD1d interaction and killer Ig receptors. [J Immunol. 181 \(5\): 3268-76.](#)
5. Hilton, H.G. *et al.* (2015) The production of KIR-Fc fusion proteins and their use in a multiplex HLA class I binding assay. [J Immunol Methods. 425: 79-87.](#)
6. Moesta, A.K. *et al.* (2009) Chimpanzees use more varied receptors and ligands than humans for inhibitory killer cell Ig-like receptor recognition of the MHC-C1 and MHC-C2 epitopes. [J Immunol. 182 \(6\): 3628-37.](#)
7. Van Der Ploeg, K. *et al.* (2017) Modulation of Human Leukocyte Antigen-C by Human Cytomegalovirus Stimulates KIR2DS1 Recognition by Natural Killer Cells. [Front Immunol. 8: 298.](#)
8. Wang, Y. *et al.* (2009) Characteristics of expanded CD4+CD28null T cells in patients with chronic hepatitis B. [Immunol Invest. 38 \(5\): 434-46.](#)

Storage Prior to reconstitution store at +4°C. Following reconstitution store at +4°C.

DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life	12 months from date of reconstitution.
-------------------	--

Health And Safety Information	Material Safety Datasheet documentation #10075 available at: 10075: https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:RPE \(MCA928PE\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M343108:190110'

Printed on 10 Jan 2019

© 2019 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)