

## Datasheet: MCA2244

<b>Description:</b>	RAT ANTI MOUSE LYMPHOTOXIN BETA RECEPTOR
<b>Specificity:</b>	LYMPHOTOXIN BETA RECEPTOR
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	5G11b
<b>Isotype:</b>	IgG2a
<b>Quantity:</b>	0.25 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Mouse
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide
<b>Carrier Free</b>	Yes

<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	Recombinant extracellular domain of LT beta R.
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P50284</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">17000</a> Ltbr    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	Tnfr, Tnfrsf3
<b>RRID</b>	AB_566581
<b>Fusion Partners</b>	Spleen cells from immunised Wistar rats were fused with cells of the SP2/0 - Ag14 mouse myeloma cell line.
<b>Specificity</b>	<p><b>Rat anti Mouse Lymphotoxin beta Receptor antibody, clone 5G11b</b> recognizes murine Tumor necrosis factor receptor superfamily member 3, also known as lymphotoxin-beta receptor, LT beta R, or Tumor necrosis factor receptor superfamily member 3. Lymphotoxin beta receptor is 385 amino acid (415 aa including 30 aa signal peptide) ~50 kDa type 1, single pass transmembrane glycoprotein and member of the tumor necrosis factor family. LT beta R is primarily expressed on stromal cells and fibroblasts and at lower levels on myeloid cell lines. Rat anti Mouse Lymphotoxin beta Receptor antibody, clone 5G11b recognizes surface expressed LT beta R.</p> <p>Rat anti Mouse Lymphotoxin beta Receptor antibody, clone 5G11b is reported to be agonistic with respect to NF Kappa B activation, MIP-2 induction and IL-6 release (<a href="#">Hehlgans et al. 2003</a>).</p> <p>Rat anti Mouse Lymphotoxin beta Receptor antibody, clone 5G11b is routinely tested on WEH164S cells.</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
<b>References</b>	<ol style="list-style-type: none"> <li>1. Palumbo, M.O. <i>et al.</i> (2006) Isolation and characterization of proinsulin-producing medullary thymic epithelial cell clones. <a href="#">Diabetes. 55: 2595-601.</a></li> <li>2. Moore, F. <i>et al.</i> (2007) An alternative pathway of NF-kappaB activation results in maturation and T cell priming activity of dendritic cells overexpressing a mutated IkappaBalpha. <a href="#">J Immunol. 178: 1301-11.</a></li> <li>3. Magari, M. <i>et al.</i> (2011) IL-21-Dependent B Cell Death Driven by Prostaglandin E2, a Product Secreted from Follicular Dendritic Cells. <a href="#">J Immunol. 187: 4210-8.</a></li> <li>4. Into, T. <i>et al.</i> (2018) MyD88 signaling causes autoimmune sialadenitis through formation of high endothelial venules and upregulation of LTββ receptor-mediated signaling. <a href="#">Sci Rep. 8 (1): 14272.</a></li> </ol>

**Further Reading** 1. Muller, P. *et al.* (2001) Functional characterization of the mouse lymphotoxin-beta receptor promoter. [Eur Cytokine Netw. 12 \(2\): 325-30.](#)

**Storage** This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

**Guarantee** 12 months from date of despatch

**Health And Safety Information** Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

**Regulatory** For research purposes only

## Related Products

### Recommended Secondary Antibodies

Goat Anti Rat IgG (STAR69...)	<a href="#">FITC</a>
Goat Anti Rat IgG (STAR73...)	<a href="#">RPE</a>
Rabbit Anti Rat IgG (STAR16...)	<a href="#">DyLight®800</a>
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	<a href="#">DyLight®800</a>
Goat Anti Rat IgG (STAR72...)	<a href="#">HRP</a>
Rabbit Anti Rat IgG (STAR21...)	<a href="#">HRP</a>
Rabbit Anti Rat IgG (STAR17...)	<a href="#">FITC</a>
Goat Anti Rat IgG (STAR131...)	<a href="#">Alk. Phos.</a> , <a href="#">Biotin</a>

### Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL \(MCA1212\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M383746:210513'

Printed on 19 May 2022

© 2022 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)