

## Datasheet: MCA2244EL

<b>Description:</b>	RAT ANTI MOUSE LYMPHOTOXIN BETA RECEPTOR:Low Endotoxin
<b>Specificity:</b>	LYMPHOTOXIN BETA RECEPTOR
<b>Format:</b>	Low Endotoxin
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	5G11b
<b>Isotype:</b>	IgG2a
<b>Quantity:</b>	0.5 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	
Functional Assays	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Mouse
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	None present
<b>Carrier Free</b>	Yes

<b>Endotoxin Level</b>	< 0.01 EU/μg
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	Recombinant extracellular domain of LT beta R.
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P50284</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">17000</a> Ltbr    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	Tnfr, Tnfrsf3
<b>RRID</b>	AB_566585
<b>Fusion Partners</b>	Spleen cells from immunized Wistar rats were fused with cells of the SP2/0 - Ag14 mouse myeloma cell line.
<b>Specificity</b>	<p><b>Rat anti Mouse Lymphotoxin beta Receptor antibody, clone 5G11b</b> recognizes murine Tumor necrosis factor receptor superfamily member 3, also known as lymphotoxin-beta receptor, LT beta R, or Tumor necrosis factor receptor superfamily member 3. Lymphotoxin beta receptor is 385 amino acid (415 aa including 30 aa signal peptide) ~50 kDa type 1, single pass transmembrane glycoprotein and member of the tumor necrosis factor family. LT beta R is primarily expressed on stromal cells and fibroblasts and at lower levels on myeloid cell lines. Rat anti Mouse Lymphotoxin beta Receptor antibody, clone 5G11b recognizes surface expressed LT beta R.</p> <p>Rat anti Mouse Lymphotoxin beta Receptor antibody, clone 5G11b is reported to be agonistic with respect to NF Kappa B activation, MIP-2 induction and IL-6 release (<a href="#">Hehlgans <i>et al.</i> 2003</a>).</p> <p>Rat anti Mouse Lymphotoxin beta Receptor antibody, clone 5G11b is routinely tested on WEH164S cells.</p>
<b>Flow Cytometry</b>	Use 10μl of the suggested working dilution to label 10 <sup>6</sup> cells in 100μl
<b>References</b>	<ol style="list-style-type: none"> <li>Palumbo, M.O. <i>et al.</i> (2006) Isolation and characterization of proinsulin-producing medullary thymic epithelial cell clones. <a href="#">Diabetes. 55: 2595-601.</a></li> <li>Moore, F. <i>et al.</i> (2007) An alternative pathway of NF-kappaB activation results in maturation and T cell priming activity of dendritic cells overexpressing a mutated IkappaBalpha. <a href="#">J Immunol. 178: 1301-11.</a></li> <li>Magari, M. <i>et al.</i> (2011) IL-21-Dependent B Cell Death Driven by Prostaglandin E2, a Product Secreted from Follicular Dendritic Cells. <a href="#">J Immunol. 187: 4210-8.</a></li> <li>Into, T. <i>et al.</i> (2018) MyD88 signaling causes autoimmune sialadenitis through formation of high endothelial venules and upregulation of LTββ receptor-mediated signaling. <a href="#">Sci</a></li> </ol>

---

<b>Further Reading</b>	1. Muller, P. <i>et al.</i> (2001) Functional characterization of the mouse lymphotoxin-beta receptor promoter. <a href="#">Eur Cytokine Netw. 12 (2): 325-30.</a>
<b>Storage</b>	Store at -20°C only. This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
<b>Guarantee</b>	12 months from date of despatch
<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10162 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA2244EL">https://www.bio-rad-antibodies.com/SDS/MCA2244EL</a>
<b>Regulatory</b>	For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16...)	<a href="#">DyLight@800</a>
Rabbit Anti Rat IgG (STAR17...)	<a href="#">FITC</a>
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	<a href="#">DyLight@550</a> , <a href="#">DyLight@650</a> , <a href="#">DyLight@800</a>
Goat Anti Rat IgG (STAR69...)	<a href="#">FITC</a>
Goat Anti Rat IgG (STAR73...)	<a href="#">RPE</a>
Goat Anti Rat IgG (STAR72...)	<a href="#">HRP</a>
Goat Anti Rat IgG (STAR131...)	<a href="#">Alk. Phos.</a> , <a href="#">Biotin</a>
Rabbit Anti Rat IgG (STAR21...)	<a href="#">HRP</a>

### Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL:Low Endotoxin \(MCA1212EL\)](#)

**Product inquiries:** [www.bio-rad-antibodies.com/technical-support](http://www.bio-rad-antibodies.com/technical-support)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](http://bio-rad-antibodies.com/datasheets)

'M445888:250929'

Printed on 06 May 2026