

Datasheet: MCA2244GA

BATCH NUMBER 1706

Description: RAT ANTI MOUSE LYMPHOTOXIN BETA RECEPTOR Specificity: LYMPHOTOXIN BETA RECEPTOR Format: Purified Product Type: Monoclonal Antibody Clone: 5G11b Isotype: IgG2a Quantity: 0.1 mg				
Format: Purified Product Type: Monoclonal Antibody Clone: 5G11b Isotype: IgG2a	Description:	RAT ANTI MOUSE LYMPHOTOXIN BETA RECEPTOR		
Product Type: Monoclonal Antibody Clone: 5G11b Isotype: IgG2a	Specificity:	LYMPHOTOXIN BETA RECEPTOR		
Clone: 5G11b Isotype: IgG2a	Format:	Purified		
Isotype: IgG2a	Product Type:	Monoclonal Antibody		
	Clone:	5G11b		
Quantity: 0.1 mg	Isotype:	lgG2a		
	Quantity:	0.1 mg		

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			1/50 - 1/100
Immunohistology - Frozen				
Immunohistology - Paraffin				
ELISA				
Immunoprecipitation				
Western Blotting				

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse	
Product Form	Purified IgG - liquid	
Preparation	Purified IgG prepared by affinity chromatography on Protein G supernatant	from tissue culture
Buffer Solution	Phosphate buffered saline	
Preservative Stabilisers	0.09% Sodium Azide	
Carrier Free	Yes	

Approx. Protein Concentrations	IgG concentration 1.0 mg/ml		
Immunogen	Recombinant extracellular domain of LT beta R.		
External Database Links	UniProt: P50284 Related reagents		
	Entrez Gene: 17000 Ltbr Related reagents		
Synonyms	Tnfcr, Tnfrsf3		
RRID	AB_566582		
Fusion Partners	Spleen cells from immunised Wistar rats were fused with cells of the SP2/0 - Ag14 mouse myeloma cell line.		
Specificity	Rat anti Mouse Lymphotoxin beta Receptor antibody, clone 5G11b recognizes murine Tumor necrosis factor receptor superfamily member 3, also known as lymphotoxin-beta receptor, LT beta R, or Tumor necrosis factor receptor superfamily member 3. Lymphotoxin beta receptor is 385 amino acid (415 aa including 30 aa signal peptide) ~50 kDa type 1, single pass transmembrane glycoprotein and member of the tumor necrosis factor family. LT beta R is primarily expressed on stromal cells and fibroblasts and at lower levels on myeloid cell lines. Rat anti Mouse Lymphotoxin beta Receptor antibody, clone 5G11b recognizes surface expressed LT beta R.		
	agonistic with respect to NF Kappa B activation, MIP-2 induction and IL-6 release (Hehlgans <i>et al.</i> 2003).		
	Rat anti Mouse Lymphotoxin beta Receptor antibody, clone 5G11b is routinely tested on WEH164S cells.		
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.		
References	 Palumbo, M.O. <i>et al.</i> (2006) Isolation and characterization of proinsulin-producing medullary thymic epithelial cell clones. <u>Diabetes. 55: 2595-601.</u> Moore, F. <i>et al.</i> (2007) An alternative pathway of NF-kappaB activation results in maturation and T cell priming activity of dendritic cells overexpressing a mutated IkappaBalpha. <u>J Immunol. 178: 1301-11.</u> Magari, M. <i>et al.</i> (2011) IL-21-Dependent B Cell Death Driven by Prostaglandin E2, a Product Secreted from Follicular Dendritic Cells. <u>J Immunol. 187: 4210-8.</u> Into, T. <i>et al.</i> (2018) MyD88 signaling causes autoimmune sialadenitis through formation of high endothelial venules and upregulation of LTββ receptor-mediated signaling. <u>Sci Rep. 8 (1): 14272.</u> 		

Further Reading	1. Muller, P. <i>et al.</i> (2001) Functional characterization of the mouse lymphotoxin-beta receptor promoter. <u>Eur Cytokine Netw. 12 (2): 325-30.</u>
Storage	Store at +4°C or at -20°C if preferred.
	This product should be stored undiluted.
	Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA2244GA 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16...)

Rabbit Anti Rat IgG (STAR17...)

Goat Anti Rat IgG (STAR72...)

HRP

Goat Anti Rat IgG (STAR69...)

Goat Anti Rat IgG (STAR73...)

Rabbit Anti Rat IgG (STAR21...)

HRP

Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...) <u>DyLight®550</u>, <u>DyLight®650</u>, <u>DyLight®800</u>

Goat Anti Rat IgG (STAR131...) Alk. Phos., Biotin

Recommended Negative Controls

RAT IgG2a NEGATIVE CONTROL (MCA1212)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M366433:200529'

Printed on 26 Jun 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint