

Datasheet: MCA2245B

BATCH NUMBER 155277

| | |
|----------------------|----------------------------|
| Description: | RAT ANTI MOUSE CD41:Biotin |
| Specificity: | CD41 |
| Other names: | INTEGRIN ALPHA IIB |
| Format: | Biotin |
| Product Type: | Monoclonal Antibody |
| Clone: | MWReg30 |
| Isotype: | IgG1 |
| Quantity: | 0.1 mg |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|----------------|-----|----|----------------|--------------------|
| Flow Cytometry | ▪ | | | Neat - 1/10 |

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

| | |
|---------------------------------------|---|
| Target Species | Mouse |
| Product Form | Purified IgG conjugated to Biotin - liquid |
| Preparation | Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant |
| Buffer Solution | Phosphate buffered saline |
| Preservative | 0.09% Sodium Azide |
| Stabilisers | 1% Bovine Serum Albumin |
| Approx. Protein Concentrations | IgG concentration 0.1 mg/ml |
| Immunogen | Purified murine platelets |

External Database**Links****UniProt:**[Q9QUM0](#)[Related reagents](#)**Entrez Gene:**[16399](#)

Itga2b

[Related reagents](#)

RRID

AB_323941

Specificity

Rat anti Mouse CD41 antibody, clone MWReg30 recognizes the mouse integrin alpha IIb subunit CD41. CD41 is a ~125 kDa single pass type 1 transmembrane glycoprotein expressed by platelets, megakaryocytes ([Zhang et al. 2007](#)), mast cells ([Berlanga et al. 2005](#)), and hematopoietic progenitors ([Mitjavila-Garcia et al. 2002](#)). CD41 forms a heterodimer with [CD61](#).

The CD41/CD61 complex is important for platelet adhesion and aggregation ([Patel et al. 2003](#)) acting as a receptor for many extracellular matrix proteins including fibronectin, thrombospondin and vitronectin ([Weisel et al. 1992](#)).

Rat anti mouse CD41, clone MWReg30 has been reported to inhibit PMA induced aggregation *in vitro* and to induce hypothermia *in vivo* ([Nieswandt et al. 1999](#)).

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity fc receptors. This may be reduced by using SeroBlock FcR ([BUF041A/B](#)).

References

1. Winter, O. *et al.* (2010) Megakaryocytes constitute a functional component of a plasma cell niche in the bone marrow. [Blood. 116: 1867-75.](#)
2. Tamagawa-Mineoka, R. *et al.* (2007) The role of platelets in leukocyte recruitment in chronic contact hypersensitivity induced by repeated elicitation. [Am J Pathol. 170: 2019-29.](#)
3. Takayama, M. *et al.* (2010) Genetic analysis of hierarchical regulation for Gata1 and NF-E2 p45 gene expression in megakaryopoiesis. [Mol Cell Biol. 30: 2668-80.](#)
4. Larson, M.K. and Watson, S.P. (2006) Regulation of proplatelet formation and platelet release by integrin alpha IIb beta3. [Blood. 108: 1509-14.](#)
5. Zanzinger, K. *et al.* (2009) Regulation of triggering receptor expressed on myeloid cells 1 expression on mouse inflammatory monocytes. [Immunology. 128: 185-95.](#)
6. Lutskiy, M.I. *et al.* (2007) WASP localizes to the membrane skeleton of platelets. [Br J Haematol. 139: 98-105.](#)
7. Sullivan, B.P. *et al.* (2010) Protective and damaging effects of platelets in acute cholestatic liver injury revealed by depletion and inhibition strategies. [Toxicol Sci. 115: 286-94.](#)
8. Fujita, R. *et al.* (2013) NF-E2 p45 Is Important for Establishing Normal Function of Platelets. [Mol Cell Biol. 33: 2659-70.](#)
9. Perez, L.E. *et al.* (2008) SH2-inositol phosphatase 1 negatively influences early megakaryocyte progenitors. [PLoS One. 3: e3565.](#)
10. Teeling, J.L. *et al.* (2012) Intracerebral immune complex formation induces

inflammation in the brain that depends on Fc receptor interaction [Acta Neuropathol. 124: 479-90.](#)

11. Motohashi, H. *et al.* (2010) NF-E2 domination over Nrf2 promotes ROS accumulation and megakaryocytic maturation. [Blood. 115 \(3\): 677-86.](#)

12. Flierl, U. *et al.* (2015) Phosphorothioate backbone modifications of nucleotide-based drugs are potent platelet activators. [J Exp Med. 212 \(2\): 129-37.](#)

13. Devanathan, V. *et al.* (2015) Platelet Gi protein Gai2 is an essential mediator of thrombo-inflammatory organ damage in mice. [Proc Natl Acad Sci U S A. 112 \(20\): 6491-6.](#)

14. Woods, S.J. *et al.* (2015) Kinetic profiling of *in vivo* lung cellular inflammatory responses to mechanical ventilation. [Am J Physiol Lung Cell Mol Physiol. 308 \(9\): L912-21.](#)

15. Goggs, R. *et al.* (2013) The small GTPase Rif is dispensable for platelet filopodia generation in mice. [PLoS One. 8 \(1\): e54663.](#)

16. Williams, C.M. *et al.* (2016) Identification of roles for the SNARE-associated protein, SNAP29, in mouse platelets. [Platelets. 27 \(4\): 286-94.](#)

17. Cuccurullo, A. *et al.* (2016) Blockade of Thrombopoietin Reduces Organ Damage in Experimental Endotoxemia and Polymicrobial Sepsis. [PLoS One. 11 \(3\): e0151088.](#)

18. Criel, M. *et al.* (2016) Absence of Pear1 does not affect murine platelet function *in vivo*. [Thromb Res. 146: 76-83.](#)

19. Ryan, J. *et al.* (2016) Myeloid cell-mediated renal injury in rapidly progressive glomerulonephritis depends upon spleen tyrosine kinase. [J Pathol. 238 \(1\): 10-20.](#)

20. Thomson, A.K. *et al.* (2017) Survival of motor neurone protein is required for normal postnatal development of the spleen. [J Anat. 230 \(2\): 337-46.](#)

21. Asai, J. *et al.* (2016) Platelets Regulate the Migration of Keratinocytes via Podoplanin/CLEC-2 Signaling during Cutaneous Wound Healing in Mice. [Am J Pathol. 186 \(1\): 101-8.](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #10041 available at: <https://www.bio-rad-antibodies.com/SDS/MCA2245B>
10041

Regulatory

For research purposes only

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M366438:200529'

