

Datasheet: MCA2245GA

BATCH NUMBER 171509

Description:	RAT ANTI MOUSE CD41
Specificity:	CD41
Other names:	INTEGRIN ALPHA IIB
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	MWReg30
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% sodium azide (NaN ₃)

Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Purified murine platelets
External Database Links	<p>UniProt: Q9QUM0 Related reagents</p> <p>Entrez Gene: 16399 Itga2b Related reagents</p>
RRID	AB_324398
Specificity	<p>Rat anti Mouse CD41 antibody, clone MWReg30 recognizes the mouse integrin alpha IIb subunit CD41. CD41 is a ~125 kDa single pass type 1 transmembrane glycoprotein expressed by platelets, megakaryocytes (Zhang <i>et al.</i> 2007), mast cells (Berlanga <i>et al.</i> 2005), and hematopoietic progenitors (Mitjavila-Garcia <i>et al.</i> 2002). CD41 forms a heterodimer with CD61.</p> <p>The CD41/CD61 complex is important for platelet adhesion and aggregation (Patel <i>et al.</i> 2003) acting as a receptor for many extracellular matrix proteins including fibronectin, thrombospondin and vitronectin (Weisel <i>et al.</i> 1992).</p> <p>Rat anti mouse CD41, clone MWReg30 has been reported to inhibit PMA induced aggregation <i>in vitro</i> and to induce hypothermia <i>in vivo</i> (Nieswandt <i>et al.</i> 1999).</p>
Flow Cytometry	Use 10µl of the suggested working dilution to label 10 ⁶ cells in 100µl
References	<ol style="list-style-type: none"> Larson, M.K. and Watson, S.P. (2006) Regulation of proplatelet formation and platelet release by integrin alpha IIb beta3. Blood. 108: 1509-14. Tamagawa-Mineoka, R. <i>et al.</i> (2007) The role of platelets in leukocyte recruitment in chronic contact hypersensitivity induced by repeated elicitation. Am J Pathol. 170: 2019-29. Lutskiy, M.I. <i>et al.</i> (2007) WASP localizes to the membrane skeleton of platelets. Br J Haematol. 139: 98-105. Perez, L.E. <i>et al.</i> (2008) SH2-inositol phosphatase 1 negatively influences early megakaryocyte progenitors. PLoS One. 3: e3565. Zanzinger, K. <i>et al.</i> (2009) Regulation of triggering receptor expressed on myeloid cells 1 expression on mouse inflammatory monocytes. Immunology. 128: 185-95. Winter, O. <i>et al.</i> (2010) Megakaryocytes constitute a functional component of a plasma cell niche in the bone marrow. Blood. 116: 1867-75. Takayama, M. <i>et al.</i> (2010) Genetic analysis of hierarchical regulation for Gata1 and NF-E2 p45 gene expression in megakaryopoiesis. Mol Cell Biol. 30: 2668-80. Sullivan, B.P. <i>et al.</i> (2010) Protective and damaging effects of platelets in acute cholestatic liver injury revealed by depletion and inhibition strategies. Toxicol Sci. 115:

[286-94.](#)

9. Motohashi, H. *et al.* (2010) NF-E2 domination over Nrf2 promotes ROS accumulation and megakaryocytic maturation. [Blood. 115 \(3\): 677-86.](#)

10. Göçmen, A.Y. *et al.* (2011) Effect of resveratrol on platelet activation in hypercholesterolemic rats: CD40-CD40L system as a potential target. [Appl Physiol Nutr Metab. 36 \(3\): 323-30.](#)

11. Teeling, J.L. *et al.* (2012) Intracerebral immune complex formation induces inflammation in the brain that depends on Fc receptor interaction [Acta Neuropathol. 124: 479-90.](#)

12. Fujita, R. *et al.* (2013) NF-E2 p45 Is Important for Establishing Normal Function of Platelets. [Mol Cell Biol. 33: 2659-70.](#)

13. Goggs, R. *et al.* (2013) The small GTPase Rif is dispensable for platelet filopodia generation in mice. [PLoS One. 8 \(1\): e54663.](#)

14. Woods, S.J. *et al.* (2015) Kinetic profiling of *in vivo* lung cellular inflammatory responses to mechanical ventilation. [Am J Physiol Lung Cell Mol Physiol. 308 \(9\): L912-21.](#)

15. Flierl, U. *et al.* (2015) Phosphorothioate backbone modifications of nucleotide-based drugs are potent platelet activators. [J Exp Med. 212 \(2\): 129-37.](#)

16. Devanathan, V. *et al.* (2015) Platelet Gi protein Gai2 is an essential mediator of thrombo-inflammatory organ damage in mice. [Proc Natl Acad Sci U S A. 112 \(20\): 6491-6.](#)

17. Cuccurullo, A. *et al.* (2016) Blockade of Thrombopoietin Reduces Organ Damage in Experimental Endotoxemia and Polymicrobial Sepsis. [PLoS One. 11 \(3\): e0151088.](#)

18. Ryan, J. *et al.* (2016) Myeloid cell-mediated renal injury in rapidly progressive glomerulonephritis depends upon spleen tyrosine kinase. [J Pathol. 238 \(1\): 10-20.](#)

19. Criel, M. *et al.* (2016) Absence of Pear1 does not affect murine platelet function *in vivo*. [Thromb Res. 146: 76-83.](#)

20. Asai, J. *et al.* (2016) Platelets Regulate the Migration of Keratinocytes via Podoplanin/CLEC-2 Signaling during Cutaneous Wound Healing in Mice. [Am J Pathol. 186 \(1\): 101-8.](#)

21. Williams, C.M. *et al.* (2016) Identification of roles for the SNARE-associated protein, SNAP29, in mouse platelets. [Platelets. 27 \(4\): 286-94.](#)

22. Thomson, A.K. *et al.* (2017) Survival of motor neurone protein is required for normal postnatal development of the spleen. [J Anat. 230 \(2\): 337-46.](#)

23. Moore, S.F. *et al.* (2021) Opposing Roles of GSK3 α and GSK3 β Phosphorylation in Platelet Function and Thrombosis. [Int J Mol Sci. 22\(19\):10656.](#)

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA2245GA>

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16...)	DyLight@800
Rabbit Anti Rat IgG (STAR17...)	FITC
Rabbit Anti Rat IgG (STAR21...)	HRP
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight@550 , DyLight@650 , DyLight@800
Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR73...)	RPE
Goat Anti Rat IgG (STAR72...)	HRP
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M413532:221123'

Printed on 23 Jan 2026