

Datasheet: MCA2245SBV570 BATCH NUMBER 100008262

Description:	RAT ANTI MOUSE CD41:StarBright Violet 570
Specificity:	CD41
Other names:	INTEGRIN ALPHA IIB
Format:	StarBright Violet 570
Product Type:	Monoclonal Antibody
Clone:	MWReg30
Isotype:	lgG1
Quantity:	100 TESTS/0.5ml

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .					
		Yes No	Not Determined	Suggested Dilution		
	Flow Cytometry			Neat		
	necessarily exclude it	s use in such proo mmended that th	for use in a particular tec edures. Suggested working a user titrates the product tive controls.	ng dilutions are given as		
Target Species	Mouse					
Product Form	Purified IgG conjugate	ed to StarBright V	olet 570 - liquid			
Max Ex/Em	Fluorophore	Excitation Max (nm) Emission Max (nm)			
	StarBright Violet 570	404	571			
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant					
Buffer Solution	Phosphate buffered saline					
Preservative	0.09% Sodium Azide (NaN ₃)					
Stabilisers	1% Bovine Serum Albumin					
	0.1% Pluronic F68					
	0.1% PEG 3350					

	0.05% Tween 20
Immunogen	Purified murine platelets
External Database Links	UniProt: <u>Q9QUM0</u> <u>Related reagents</u> Entrez Gene: <u>16399</u> Itga2b <u>Related reagents</u>
Specificity	Rat anti Mouse CD41 antibody, clone MWReg30 recognizes the mouse integrin alpha Ilb subunit CD41. CD41 is a ~125 kDa single pass type 1 transmembrane glycoprotein expressed by platelets, megakaryocytes (Zhang <i>et al.</i> 2007), mast cells (Berlanga <i>et al.</i> 2005), and hematopoietic progenitors (Mitjavila-Garcia <i>et al.</i> 2002). CD41 forms a heterodimer with CD61. The CD41/CD61 complex is important for platelet adhesion and aggregation (Patel <i>et al.</i> 2003) acting as a receptor for many extracellular matrix proteins including fibronectin, thrombospondin and vitronectin (Weisel <i>et al.</i> 1992).
	Rat anti mouse CD41, clone MWReg30 has been reported to inhibit PMA induced aggregation <i>in vitro</i> and to induce hypothermia <i>in vivo</i> (<u>Nieswandt <i>et al.</i> 1999</u>).
Flow Cytometry	Use 5µl of the suggested working dilution to label 10 ⁶ cells in 100µl. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.
References	 Larson, M.K. and Watson, S.P. (2006) Regulation of proplatelet formation and platelet release by integrin alpha IIb beta3. <u>Blood. 108: 1509-14.</u> Tamagawa-Mineoka, R. <i>et al.</i> (2007) The role of platelets in leukocyte recruitment in chronic contact hypersensitivity induced by repeated elicitation. <u>Am J Pathol. 170: 2019-29.</u> Lutskiy, M.I. <i>et al.</i> (2007) WASP localizes to the membrane skeleton of platelets. <u>Br J Haematol. 139: 98-105.</u> Perez, L.E. <i>et al.</i> (2008) SH2-inositol phosphatase 1 negatively influences early megakaryocyte progenitors. <u>PLoS One. 3: e3565.</u> Zanzinger, K. <i>et al.</i> (2009) Regulation of triggering receptor expressed on myeloid cells 1 expression on mouse inflammatory monocytes. <u>Immunology. 128: 185-95.</u> Winter, O. <i>et al.</i> (2010) Megakaryocytes constitute a functional component of a plasma cell niche in the bone marrow. <u>Blood. 116: 1867-75.</u> Takayama, M. <i>et al.</i> (2010) Protective and damaging effects of platelets in acute cholestatic liver injury revealed by depletion and inhibition strategies. <u>Toxicol Sci. 115: 286-94.</u> Motohashi, H. <i>et al.</i> (2010) NF-E2 domination over Nrf2 promotes ROS accumulation and megakaryocytic maturation. <u>Blood. 115 (3): 677-86.</u> Göçmen, A.Y. <i>et al.</i> (2011) Effect of resveratrol on platelet activation in

	 hypercholesterolemic rats: CD40-CD40L system as a potential target. <u>Appl Physiol Nutr</u> <u>Metab. 36 (3): 323-30.</u> 11. Teeling, J.L. <i>et al.</i> (2012) Intracerebral immune complex formation induces inflammation in the brain that depends on Fc receptor interaction <u>Acta Neuropathol. 124:</u> <u>479-90.</u> 12. Eviita, P. <i>et al.</i> (2012) NE E2 p45 is important for Establishing Normal Europian of
	12. Fujita, R. <i>et al.</i> (2013) NF-E2 p45 Is Important for Establishing Normal Function of Platelets. <u>Mol Cell Biol. 33: 2659-70.</u>
	13. Goggs, R. <i>et al.</i> (2013) The small GTPase Rif is dispensable for platelet filopodia
	generation in mice. <u>PLoS One. 8 (1): e54663.</u>
	14. Woods, S.J. <i>et al.</i> (2015) Kinetic profiling of <i>in vivo</i> lung cellular inflammatory
	responses to mechanical ventilation. Am J Physiol Lung Cell Mol Physiol. 308 (9):
	<u>L912-21.</u> 15. Flight LL et el. (2015) Pheenherethieste heekhene medifications of nucleotide heesd
	15. Flierl, U. <i>et al.</i> (2015) Phosphorothioate backbone modifications of nucleotide-based drugs are potent platelet activators. J Exp Med. 212 (2): 129-37.
	16. Devanathan, V. <i>et al.</i> (2015) Platelet Gi protein Gαi2 is an essential mediator of
	thrombo-inflammatory organ damage in mice. <u>Proc Natl Acad Sci U S A. 112 (20): 6491-6.</u>
	17. Cuccurullo, A. <i>et al.</i> (2016) Blockade of Thrombopoietin Reduces Organ Damage in
	Experimental Endotoxemia and Polymicrobial Sepsis. PLoS One. 11 (3): e0151088.
	18. Ryan, J. <i>et al.</i> (2016) Myeloid cell-mediated renal injury in rapidly progressive
	glomerulonephritis depends upon spleen tyrosine kinase. <u>J Pathol. 238 (1): 10-20.</u>
	19. Criel, M. <i>et al.</i> (2016) Absence of Pear1 does not affect murine platelet function <i>in vivo</i> . <u>Thromb Res. 146: 76-83.</u>
	20. Asai, J. <i>et al.</i> (2016) Platelets Regulate the Migration of Keratinocytes via
	Podoplanin/CLEC-2 Signaling during Cutaneous Wound Healing in Mice. Am J Pathol.
	<u>186 (1): 101-8.</u>
	21. Williams, C.M. <i>et al.</i> (2016) Identification of roles for the SNARE-associated protein, SNAP29, in mouse platelets. <u>Platelets. 27 (4): 286-94.</u>
	22. Thomson, A.K. et al. (2017) Survival of motor neurone protein is required for normal
	postnatal development of the spleen. <u>J Anat. 230 (2): 337-46.</u>
	23. Moore, S.F. <i>et al.</i> (2021) Opposing Roles of GSK3α and GSK3β Phosphorylation in
	Platelet Function and Thrombosis. Int J Mol Sci. 22(19):10656.
Storage	Store at +4°C. DO NOT FREEZE.
	This product should be stored undiluted.
Guarantee	12 months from date of despatch
Acknowledgements	This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign counterparts
Health And Safety Information	Material Safety Datasheet documentation #20471 available at: https://www.bio-rad-antibodies.com/SDS/MCA2245SBV570 20471
Regulatory	For research purposes only

Related Products

Recommended Useful Reagents

MOUSE SEROBLOCK FcR (BUF041A) MOUSE SEROBLOCK FcR (BUF041B)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@b	io-rad.com	Email: antibody_sales_uk@bio	-rad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M426137:231121'

Printed on 29 Apr 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint