

Datasheet: MCA2245SBV570

Description:	RAT ANTI MOUSE CD41:StarBright Violet 570
Specificity:	CD41
Other names:	INTEGRIN ALPHA IIB
Format:	StarBright Violet 570
Product Type:	Monoclonal Antibody
Clone:	MWReg30
Isotype:	IgG1
Quantity:	100 TESTS/0.5ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse				
Product Form	Purified IgG conjugat	Purified IgG conjugated to StarBright Violet 570 - liquid			
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm		
	StarBright Violet 570	404	571		
5					
Preparation	Purified IgG prepared supernatant	d by affinity chromatog	raphy on Protein G		
Buffer Solution	•		raphy on Protein G		
	supernatant	saline	raphy on Protein G		
Buffer Solution	supernatant Phosphate buffered s	caline (NaN ₃)	raphy on Protein G		
Buffer Solution Preservative	Phosphate buffered s	caline (NaN ₃)	raphy on Protein G		
Buffer Solution Preservative	Supernatant Phosphate buffered s 0.09% Sodium Azide 1% Bovine Serum All	caline (NaN ₃)	raphy on Protein G		

lm	m	 nn	a	n

Purified murine platelets

External Database Links

UniProt:

Q9QUM0 Related reagents

Entrez Gene:

16399 Itga2b Related reagents

Specificity

Rat anti Mouse CD41 antibody, clone MWReg30 recognizes the mouse integrin alpha IIb subunit CD41. CD41 is a ~125 kDa single pass type 1 transmembrane glycoprotein expressed by platelets, megakaryocytes (Zhang et al. 2007), mast cells (Berlanga et al. 2005), and hematopoietic progenitors (Mitjavila-Garcia et al. 2002). CD41 forms a heterodimer with CD61.

The CD41/CD61 complex is important for platelet adhesion and aggregation (<u>Patel et al. 2003</u>) acting as a receptor for many extracellular matrix proteins including fibronectin, thrombospondin and vitronectin (<u>Weisel et al. 1992</u>).

Rat anti mouse CD41, clone MWReg30 has been reported to inhibit PMA induced aggregation *in vitro* and to induce hypothermia *in vivo* (Nieswandt *et al.* 1999).

Flow Cytometry

Use 5μ I of the suggested working dilution to label 10^6 cells in 100μ I. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.

References

- 1. Larson, M.K. and Watson, S.P. (2006) Regulation of proplatelet formation and platelet release by integrin alpha IIb beta3. Blood. 108: 1509-14.
- 2. Tamagawa-Mineoka, R. *et al.* (2007) The role of platelets in leukocyte recruitment in chronic contact hypersensitivity induced by repeated elicitation. <u>Am J Pathol. 170:</u> 2019-29.
- 3. Lutskiy, M.I. *et al.* (2007) WASP localizes to the membrane skeleton of platelets. <u>Br J Haematol</u>. 139: 98-105.
- 4. Perez, L.E. *et al.* (2008) SH2-inositol phosphatase 1 negatively influences early megakaryocyte progenitors. <u>PLoS One. 3: e3565.</u>
- 5. Zanzinger, K. *et al.* (2009) Regulation of triggering receptor expressed on myeloid cells 1 expression on mouse inflammatory monocytes. Immunology. 128: 185-95.
- 6. Winter, O. *et al.* (2010) Megakaryocytes constitute a functional component of a plasma cell niche in the bone marrow. <u>Blood.</u> 116: 1867-75.
- 7. Takayama, M. *et al.* (2010) Genetic analysis of hierarchical regulation for Gata1 and NF-E2 p45 gene expression in megakaryopoiesis. Mol Cell Biol. 30: 2668-80.
- 8. Sullivan, B.P. *et al.* (2010) Protective and damaging effects of platelets in acute cholestatic liver injury revealed by depletion and inhibition strategies. <u>Toxicol Sci. 115:</u> <u>286-94.</u>
- 9. Motohashi, H. *et al.* (2010) NF-E2 domination over Nrf2 promotes ROS accumulation and megakaryocytic maturation. <u>Blood. 115 (3): 677-86.</u>
- 10. Göçmen, A.Y. *et al.* (2011) Effect of resveratrol on platelet activation in hypercholesterolemic rats: CD40-CD40L system as a potential target. <u>Appl Physiol Nutr Metab.</u> 36 (3): 323-30.

- 11. Teeling, J.L. *et al.* (2012) Intracerebral immune complex formation induces inflammation in the brain that depends on Fc receptor interaction <u>Acta Neuropathol. 124:</u> 479-90.
- 12. Fujita, R. *et al.* (2013) NF-E2 p45 Is Important for Establishing Normal Function of Platelets. Mol Cell Biol. 33: 2659-70.
- 13. Goggs, R. *et al.* (2013) The small GTPase Rif is dispensable for platelet filopodia generation in mice. <u>PLoS One. 8 (1): e54663.</u>
- 14. Woods, S.J. *et al.* (2015) Kinetic profiling of *in vivo* lung cellular inflammatory responses to mechanical ventilation. <u>Am J Physiol Lung Cell Mol Physiol. 308 (9):</u> L912-21.
- 15. Flierl, U. *et al.* (2015) Phosphorothioate backbone modifications of nucleotide-based drugs are potent platelet activators. <u>J Exp Med. 212 (2): 129-37.</u>
- 16. Devanathan, V. *et al.* (2015) Platelet Gi protein Gαi2 is an essential mediator of thrombo-inflammatory organ damage in mice. Proc Natl Acad Sci U S A. 112 (20): 6491-6.
- 17. Cuccurullo, A. *et al.* (2016) Blockade of Thrombopoietin Reduces Organ Damage in Experimental Endotoxemia and Polymicrobial Sepsis. PLoS One. 11 (3): e0151088.
- 18. Ryan, J. *et al.* (2016) Myeloid cell-mediated renal injury in rapidly progressive glomerulonephritis depends upon spleen tyrosine kinase. J Pathol. 238 (1): 10-20.
- 19. Criel, M. *et al.* (2016) Absence of Pear1 does not affect murine platelet function *in vivo*. Thromb Res. 146: 76-83.
- 20. Asai, J. *et al.* (2016) Platelets Regulate the Migration of Keratinocytes via Podoplanin/CLEC-2 Signaling during Cutaneous Wound Healing in Mice. <u>Am J Pathol.</u> 186 (1): 101-8.
- 21. Williams, C.M. *et al.* (2016) Identification of roles for the SNARE-associated protein, SNAP29, in mouse platelets. <u>Platelets. 27 (4): 286-94.</u>
- 22. Thomson, A.K. *et al.* (2017) Survival of motor neurone protein is required for normal postnatal development of the spleen. <u>J Anat. 230 (2): 337-46.</u>
- 23. Moore, S.F. *et al.* (2021) Opposing Roles of GSK3α and GSK3β Phosphorylation in Platelet Function and Thrombosis. Int J Mol Sci. 22(19):10656.

Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted.	
Guarantee	12 months from date of despatch	
Acknowledgements	This product is covered by U.S. Patent No. 10,150,841 and related U.S. and forei counterparts	gn
Health And Safety Information	Material Safety Datasheet documentation #20471 available at: https://www.bio-rad-antibodies.com/SDS/MCA2245SBV570 20471	
Regulatory	For research purposes only	

Related Products

Recommended Useful Reagents

MOUSE SEROBLOCK FcR (BUF041A)

MOUSE SEROBLOCK FcR (BUF041B)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M426137:231121'

Printed on 29 Apr 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint