

Datasheet: MCA2250XZ

Description:	RAT ANTI HUMAN INTERLEUKIN-10:Preservative Free
Specificity:	IL-10
Format:	Preservative Free
Product Type:	Monoclonal Antibody
Clone:	JES3-12G8
Isotype:	IgG2a
Quantity:	1 mg

Product Details

RRID AB_567072

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry (1)	▪			
Immunohistology - Frozen			▪	
Immunohistology - Paraffin (2)	▪			1/100
ELISA	▪			
Immunoprecipitation			▪	
Western Blotting	▪			1/100 - 1/250
Functional Assays	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using the appropriate negative/positive controls.

(1)**Membrane permeabilisation is required for this application. Bio-Rad recommends the use of Leucoperm™ (Product Code [BUF09](#)) for this purpose.**

(2)**This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose.**

Target Species Human

Product Form Purified IgG - liquid

Preparation Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

Buffer Solution Borate buffered saline

Preservative Stabilisers None Present

Approx. Protein Concentrations IgG concentration 0.5 mg/ml

Immunogen	Human IL-10 expressed by COS cells.
External Database Links	<p>UniProt: P22301 Related reagents</p> <p>Entrez Gene: 3586 IL10 Related reagents</p>
Specificity	<p>Rat anti Human Interleukin-10 antibody, clone JES3-12G8 recognizes human Interleukin-10 (IL-10). IL-10 is expressed by activated CD8+ peripheral blood T-cells, activated T-helper CD4+ T cell clones mast cells, B cell lymphomas and LPS activated monocytes.</p> <p>Rat anti Human Interleukin-10 antibody, clone JES3-12G8 is reported to neutralize the effects of IL-10 (Abrams <i>et al.</i> 1992).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul.
Histology Positive Control Tissue	Human Tonsil
References	<ol style="list-style-type: none"> 1. Brodeur, N.D. and Spencer, J.V. (2010) Antibodies to human IL-10 neutralize ebvIL-10-mediated cytokine suppression but have no effect on cmvIL-10 activity. Virus Res. 153: 265-8. 2. Burdin, N. <i>et al.</i> (1993) Epstein-Barr virus transformation induces B lymphocytes to produce human interleukin 10. J Exp Med. 177 (2): 295-304. 3. O'farrell, A.M. <i>et al.</i> (1998) IL-10 inhibits macrophage activation and proliferation by distinct signaling mechanisms: evidence for Stat3-dependent and -independent pathways. EMBO J. 17 (4): 1006-18. 4. Abrams, J.S. <i>et al.</i> (1992) Strategies of anti-cytokine monoclonal antibody development: immunoassay of IL-10 and IL-5 in clinical samples. Immunol Rev. 127: 5-24. 5. Wakkach, A. <i>et al.</i> (2001) Differentiation of regulatory T cells 1 is induced by CD2 costimulation. J Immunol. 167: 3107-13. 6. Dybiec, M. <i>et al.</i> (2007) Photoluminescence spectroscopy of bioconjugated CdSe/ZnS quantum dots Appl. Phys. Lett. 90, 263112. 7. Ramgir, N.S. <i>et al.</i> (2007) Voltammetric Detection of Cancer Biomarkers Exemplified by Interleukin-10 and Osteopontin with Silica Nanowires J. Phys. Chem. C, 111:13981-7. 8. Zajac, A. <i>et al.</i> (2007) Protein microarrays and quantum dot probes for early cancer detection. Colloids Surf B Biointerfaces. 58: 309-14.
Further Reading	<ol style="list-style-type: none"> 1. Gomes, J.A. <i>et al.</i> (2003) Evidence that development of severe cardiomyopathy in human Chagas' disease is due to a Th1-specific immune response. Infect Immun. 71 (3): 1185-93. 2. Yssel, H. <i>et al.</i> (1992) IL-10 is produced by subsets of human CD4+ T cell clones and peripheral blood T cells. J Immunol. 149 (7): 2378-84.
Storage	<p>Store at -20°C only.</p> <p>This product should be stored undiluted.</p> <p>Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.</p>

Guarantee 18 months from date of despatch.

Health And Safety Information Material Safety Datasheet documentation #10123 available at:
10123: <https://www.bio-rad-antibodies.com/uploads/MSDS/10123.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin
Goat Anti Rat IgG (STAR73...)	RPE
Rabbit Anti Rat IgG (STAR21...)	HRP
Goat Anti Rat IgG (STAR72...)	HRP
Rabbit Anti Rat IgG (STAR16...)	DyLight®800
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight®649 , DyLight®800

North & South America Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M341584:190109'

Printed on 16 Mar 2019

© 2019 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)