

Datasheet: MCA2255

Description:	RAT ANTI HUMAN CDw210a
Specificity:	CDw210a
Other names:	IL-10R1
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	3F9
Isotype:	IgG2a
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry (1)	■			1/10 - 1/50
Immunohistology - Frozen			■	
Immunohistology - Paraffin			■	
ELISA			■	
Immunoprecipitation			■	
Western Blotting			■	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1)Due to low expression of CDw210a, sensitive staining techniques may be required.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant.
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Soluble human CDw210a

**External Database
Links**

UniProt:

[Q13651](#) [Related reagents](#)

Entrez Gene:

[3587](#) IL10RA [Related reagents](#)

Synonyms

IL10R

Fusion Partners

Spleen cells from immunised Fisher rats were fused with cells of the mouse SP2/0 myeloma cell line.

Specificity

Rat anti Human CDw210a, clone 3F9 recognizes human CDw210a, a 90-110kD glycoprotein, also known as Interleukin-10 receptor 1 (IL-10R1).

CDw210a is expressed on T cells, NK cells, monocytes and dendritic cells, although surface expression may be low.

Recent work has reported an increase in intracellular expression of CDw210a during dendritic cell maturation, but a reduction in surface CDw210a expression, see [Corinti, S. et al.](#) for details.

This clone is reported to function as a blocking antibody. Bio-Rad recommend the use of [MCA2255XZ](#) for functional studies, see [Liu, Y. et al.](#) for details.

Flow Cytometry

Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.

References

1. Liu, Y. *et al.* (1997) The EBV IL-10 homologue is a selective agonist with impaired binding to the IL-10 receptor. [J Immunol. 158 \(2\): 604-13.](#)
2. O7#39;Farrell, A.M. *et al.* (1998) IL-10 inhibits macrophage activation and proliferation by distinct signaling mechanisms: evidence for Stat3-dependent and -independent pathways. [EMBO J. 17 \(4\): 1006-18.](#)
3. Corinti, S. *et al.* (2001) Regulatory activity of autocrine IL-10 on dendritic cell functions. [J Immunol. 166 \(7\): 4312-8.](#)
4. Elbim, C. *et al.* (2001) Intracellular pool of IL-10 receptors in specific granules of human neutrophils: differential mobilization by proinflammatory mediators. [J Immunol. 166 \(8\): 5201-7.](#)
5. Fayette, J. *et al.* (1997) Human dendritic cells skew isotype switching of CD40-activated naive B cells towards IgA1 and IgA2. [J Exp Med. 185 \(11\): 1909-18.](#)
6. Goodier, M.R. & Londei, M. (2000) Lipopolysaccharide stimulates the proliferation of human CD56+CD3- NK cells: a regulatory role of monocytes and IL-10. [J Immunol. 165 \(1\): 139-47.](#)
7. Levings, M.K. *et al.* (2001) IFN-alpha and IL-10 induce the differentiation of human type 1 T regulatory cells. [J Immunol. 166 \(9\): 5530-9.](#)
8. Hamel, Y. *et al.* (2016) A unique CD8(+) T lymphocyte signature in pediatric type 1 diabetes. [J Autoimmun. 73: 54-63.](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life

18 months from date of despatch.

**Health And Safety
Information**

Material Safety Datasheet documentation #10040 available at:
10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16...)	DyLight®800
Goat Anti Rat IgG (STAR73...)	RPE
Rabbit Anti Rat IgG (STAR21...)	HRP
Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight®649 , DyLight®800
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin
Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR72...)	HRP

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751
Email: antibody_sales_us@bio-rad.com

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739
Email: antibody_sales_uk@bio-rad.com

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50
Email: antibody_sales_de@bio-rad.com

'M323505:180727'

Printed on 27 Jul 2018