

Datasheet: MCA2257B

BATCH NUMBER 1610

Description:	MOUSE ANTI HUMAN CD226:Biotin
Specificity:	CD226
Other names:	DNAM-1
Format:	Biotin
Product Type:	Monoclonal Antibody
Clone:	DX11
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10
Immunoprecipitation	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG conjugated to Biotin - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative	0.09% Sodium Azide
Stabilisers	1.0% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml

Immunogen	Human cytotoxic T lymphocyte clone
External Database Links	<p>UniProt: Q15762 Related reagents</p> <p>Entrez Gene: 10666 CD226 Related reagents</p>
Synonyms	DNAM1
RRID	AB_324795
Fusion Partners	Spleen cells from immunized BALB/c mouse were fused with cells of the Sp2/0 myeloma cell line
Specificity	<p>Mouse anti Human CD226 antibody, clone DX11 recognizes human CD226, a ~65 kDa glycoprotein, also known as DNAM1 (DNAX accessory molecule-1). CD226 is broadly expressed on T-cells, NK cells, platelets, monocytes and a subset of B cells. CD226 is also expressed by a subset of CD3 positive thymocytes.</p> <p>Mouse anti Human CD226 antibody, clone DX11 is reported to inhibit T- and NK cell mediated cytotoxicity against tumor cell targets and to block TNF alpha and IFN gamma secretion by alloantigen-specific T-cells (Kojima et al. 2003).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 1 x 10 ⁶ cells in 100ul.
References	<ol style="list-style-type: none"> 1. Shibuya, A. <i>et al.</i> (1996) DNAM-1, a novel adhesion molecule involved in the cytolytic function of T lymphocytes. Immunity. 4 (6): 573-81. 2. Shibuya, A. <i>et al.</i> (1998) Protein kinase C is involved in the regulation of both signaling and adhesion mediated by DNAX accessory molecule-1 receptor. J Immunol. 161 (4): 1671-6. 3. Kojima, H. <i>et al.</i> (2003) CD226 mediates platelet and megakaryocytic cell adhesion to vascular endothelial cells. J Biol Chem. 278 (38): 36748-53. 4. Manes, T.D. and Pober, J.S. (2011) Identification of Endothelial Cell Junctional Proteins and Lymphocyte Receptors Involved in Transendothelial Migration of Human Effector Memory CD4+ T Cells. J Immunol. 186: 1763-8. 5. Ardolino, M. <i>et al.</i> (2011) DNAM-1 ligand expression on Ag-stimulated T lymphocytes is mediated by ROS-dependent activation of DNA-damage response: relevance for NK-T cell interaction. Blood. 117: 4778-86. 6. Soriani, A. <i>et al.</i> (2009) ATM-ATR-dependent up-regulation of DNAM-1 and NKG2D ligands on multiple myeloma cells by therapeutic agents results in enhanced NK-cell susceptibility and is associated with a senescent phenotype. Blood. 113: 3503-11. 7. Matusali, G. <i>et al.</i> (2012) The Human Immunodeficiency Virus Type 1 Nef and Vpu Proteins Downregulate the Natural Killer Cell-Activating Ligand PVR. J Virol. 86: 4496-504. 8. Fionda, C. <i>et al.</i> (2015) Nitric oxide donors increase PVR/CD155 DNAM-1 ligand expression in multiple myeloma cells: role of DNA damage response activation. BMC

[Cancer. 15 \(1\): 17.](#)

9. Molfetta, R. *et al.* (2019) The Ubiquitin-proteasome pathway regulates Nectin2/CD112 expression and impairs NK cell recognition and killing. [Eur J Immunol. 49 \(6\): 873-83.](#)

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10041 available at: <https://www.bio-rad-antibodies.com/SDS/MCA2257B>
10041

Regulatory For research purposes only

Related Products

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M366461:200529'

Printed on 29 Apr 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)