# Datasheet: MCA2259 BATCH NUMBER 150955

Description:	MOUSE ANTI OVALBUMIN
Specificity:	OVALBUMIN
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	2C6
lsotype:	IgE
Quantity:	0.1 mg

## **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal				
	communications from the originators. Please refer to references indicated for further				
	information. For general protocol recommendations, please visit www.bio-				
	rad-antibodies.com/protocols.				
		Yes	No	Not Determined	Suggested Dilution
	Flow Cytometry			•	

	Flow Cytometry	-		
	Immunohistology - Frozen	•		
	Immunohistology - Paraffin	•		
	ELISA	•	1/1000 - 1/5000	
	Immunoprecipitation	•		
	Western Blotting	•		
	Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.			
Target Species	Chicken			
Product Form	Purified IgE - liquid			
Preparation	Purified IgE prepared from	m tissue culture supernatant.		
Buffer Solution	Phosphate buffered saline			
Preservative Stabilisers	0.09% Sodium Azide			
Approx. Protein Concentrations	IgE concentration 1.0 mg	/ml		

Immunogen	Ovalbumin.
RRID	AB_2285753
Fusion Partners	Spleen cells from immunized Balb/c mice were fused with cells of the mouse myeloma, P3U1.
Specificity	<b>Mouse anti Ovalbumin antibody, clone 2C6</b> recognises ovalbumin (OVA). The antibody is suitable for use as a mouse IgE standard in ELISA assays ( <u>Hamada<i>et al.</i> 2003</u> ).
References	<ol> <li>Stevens, T. <i>et al.</i> (2008) Increased transcription of immune and metabolic pathways in naive and allergic mice exposed to diesel exhaust. Toxicol Sci. 102: 359-70.</li> <li>Fairley, K.J. <i>et al.</i> (2007) Exposure to the immunosuppressant, perfluorooctanoic acid, enhances the murine IgE and airway hyperreactivity response to ovalbumin. Toxicol Sci. 97 (2): 375-83.</li> <li>Ellertsen, L.K. <i>et al.</i> (2010) Maternal allergen immunisation to prevent sensitisation in offspring: Th2-polarising adjuvants are more efficient than a Th1-polarising adjuvant in mice. BMC Immunol. 11: 8-17</li> <li>Kambayashi, T. <i>et al.</i> (2008) Indirect involvement of allergen-captured mast cells in antigen presentation. Blood. 111:1489-96.</li> <li>Paliwal, S. <i>et al.</i> (2007) Early-life psychological stress exacerbates adult mouse asthma via the hypothalamus-pituitary-adrenal axis. Am J Respir Crit Care Med. 175: 316-22.</li> <li>Chida, Y. <i>et al.</i> (2007) A small-molecule compound targeting CCR5 and CXCR3 prevents airway hyperresponsiveness and inflammation. Eur Respir J. 31: 783-9.</li> <li>Suzaki, Y. <i>et al.</i> (2010) Determinants of experimental allergic responses: interactions between allergen dose, sex and age. Scand J Immunol. 73 (6): 554-67.</li> <li>Nygaard, U.C. <i>et al.</i> (2015) Allergen-Specific Immunotherapy with Monomeric Allergoid in a Mouse Model of Atopic Dermatitis. PLoS One. 10 (8): e0135070.</li> <li>Piro B. <i>et al.</i> (2011) Development of atopic provision is associated with protection from food allergy and oral tolerance. Food Chem Toxicol. 83: 17-25.</li> <li>Sheshakova N <i>et al.</i> (2015) Allergen-Specific Immunotherapy with Monomeric Allergoid in a Mouse Model of Atopic Dermatitis. PLoS One. 10 (8): e0135070.</li> <li>Piro B. <i>et al.</i> (2011) Development of atopy by severe paramyxoviral infection in a mouse model. Ann Allergy Asthma Immunol. 105 (6): 437-443. e1.</li> <li>Garbani, M. <i>et al.</i> (2016) A distinct microbida composition is associated with protection from fo</li></ol>

	<ul> <li>Pub: US 2015/0275174 A1</li> <li>18. Andreassen, M. <i>et al.</i> (2015) Cry1Ab protein from <i>Bacillus ta</i> cry1Ab-transgenic maize exerts no adjuvant effect after airway Immunol. 81 (3): 192-200.</li> <li>19. Shin, W.<i>et al.</i> (2018) V-set and Ig domain-containing 4 (VSI F4/80<sup>+</sup> cells regulate oral antigen-specific responses in mouse.</li> <li><u>632-43.</u></li> </ul>	exposure. <u>Scand J</u> G4)-expressing hepatic
Storage	Store at +4°C or at -20°C if preferred. Avoid repeated freezing a denature the antibody.	and thawing as this may
	Storage in frost free freezers is not recommended.	
	This product should be stored undiluted. Should this product co recommend microcentrifugation before use.	ntain a precipitate we
Guarantee	12 months from date of despatch	
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA2259 10040	
Regulatory	For research purposes only	

### **Related Products**

### **Recommended Secondary Antibodies**

#### Rat Anti Mouse IgE HEAVY CHAIN (MCA419...) HRP

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-ra	ad.com	Email: antibody_sales_uk@bio-	rad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M366471:200529'

#### Printed on 19 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint