

# Datasheet: MCA2260

Description:	n: RAT ANTI RITUXIMAB		
Specificity:	RITUXIMAB		
Other names:	MabThera , Rituxan		
Format:	Purified		
Product Type:	Monoclonal Antibody		
Clone:	MB2A4		
Isotype:	lgG2a		
Quantity:	0.2 mg		

## **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal					
	communications from the originators. Please refer to references indicated for further					
	information. For general protocol recommendations, please visit <u>www.bio-</u>					
	rad-antibodies.com/protocols.					
		Yes	No	Not Determined	Suggested Dilution	
	Flow Cytometry	-			50 ug/ml	
	Immunohistology - Frozen			•		
	Immunohistology - Paraffin			•		
	ELISA	-				
	Where this product has not been tested for use in a particular technique this does not					
	necessarily exclude its us	se in such	n procedur	es. Suggested workin	ig dilutions are given as	
	a guide only. It is recomn	nended th	at the use	r titrates the product f	for use in their own	
	system using appropriate	e negative	/positive c	ontrols.		
Product Form	Purified IgG - liquid					
Preparation	Purified IgG prepared by supernatant	affinity cł	nromatogra	aphy on Protein G froi	m tissue culture	
Buffer Solution	TRIS buffered saline					
Preservative Stabilisers	<0.1% sodium azide (Na	N <sub>3</sub> )				
Approx. Protein Concentrations	IgG concentration 1.0 mg	g/ml				
Immunogen	F(ab) <sub>2</sub> fragment of Rituxi	mab				

RRID	AB_323787
Fusion Partners	Spleen cells from immunised rats were fused with cells of the NS-1 mouse myeloma cell line
Specificity	<b>Rat Anti-Rituximab Antibody, clone MB2A4,</b> is an anti-idiotypic antibody that recognizes the monoclonal antibody drug rituximab. The antibody can be used to measure the levels of rituximab and biosimilar products in bioanalytical assays. Rat Anti-Rituximab Antibody also recognizes ocrelizumab, a second-generation humanized anti-CD20 antibody that binds to an epitope on CD20 that is identical or overlapping with the rituximab epitope. Clone MB2A4 has been used in ELISA to monitor the levels of rituximab in patient serum following therapy (Cragg <i>et al.</i> 2004 and Hampson <i>et al.</i> 2010).
	Clone MB2A4 has been used to detect rituximab bound to the surface of the Raji B cell line, however detection of rituximab bound in vivo to B-CLL cells has not been demonstrated. It is possible that complement deposition on rituximab opsonised cells inhibits binding of the Anti-Rituximab Antibody to cell bound rituximab ( <u>Beum <i>et al.</i> 2004</u> ). Inhibition experiments carried out with Daudi cells demonstrated that this antibody is inhibitory at a ratio of 5:1 antibody:rituximab, but does not inhibit rituximab binding to CD20 at a ratio of 1:1.
	Rituximab (reference product branded as Rituxan) is a chimeric mouse/human monoclonal antibody approved for the treatment of certain autoimmune diseases and cancer, including non-Hodgkin's lymphoma, chronic lymphocytic leukemia and rheumatoid arthritis. The antibody is specific for the cell surface protein CD20, which is widely expressed on B cells. Through three different mechanisms of action it eliminates B cells from the body, enabling the development of a new population of healthy B cells.
	View a summary of all Anti-Rituximab Antibodies.
Flow Cytometry	Use 10 $\mu$ I of the suggested working dilution to label 10 <sup>6</sup> cells in 100 $\mu$ I
ELISA	Rat anti rituximab antibody, clone MB2A4 may be used in a direct ELISA or as a detection reagent, when conjugated to HRP, in a sandwich ELISA together with <u>HCA186</u> as the capture reagent. Protocol: <u>PK bridging ELISA to measure free drug</u>
References	<ol> <li>Cragg, M. S. <i>et al.</i> (2004) A new anti-idiotype antibody capable of binding rituximab on the surface of lymphoma cells. <u>Blood. 104:2540-2</u></li> <li>Cragg, M.S. <i>et al.</i> (2004) Apparent modulation of CD20 by rituximab: an alternative explanation. <u>Blood. 103 (10): 3989-90; author reply 3990-1.</u></li> <li>Pers, J.O. <i>et al.</i> (2007) BAFF-modulated repopulation of B lymphocytes in the blood and salivary glands of rituximab-treated patients with Sjögren's syndrome. <u>Arthritis Rheum. 56: 1464-77.</u></li> <li>Hampson, G. <i>et al.</i> (2010) Validation of an ELISA for the determination of rituximab pharmacokinetics in clinical trials subjects. <u>J Immunol Methods. 360 (1-2): 30-8.</u></li> <li>Blasco, H. <i>et al.</i> (2007) Evaluation of a peptide ELISA for the detection of rituximab in serum. <u>J Immunol Methods.325: 127-39.</u></li> <li>Daydé, D. <i>et al.</i> (2009) Tumor burden influences exposure and response to rituximab:</li> </ol>

pharmacokinetic-pharmacodynamic modeling using a syngeneic bioluminescent murine model expressing human CD20. <u>Blood. 113: 3765-72.</u>

7. Aung, T. *et al.* (2011) Exosomal evasion of humoral immunotherapy in aggressive B-cell lymphoma modulated by ATP-binding cassette transporter A3. <u>Proc Natl Acad Sci U S A.</u> <u>108: 15336-41.</u>

8. Schmidt, E. *et al.* (2009) Immunogenicity of rituximab in patients with severe pemphigus. <u>Clin Immunol. 132: 334-41.</u>

 McDonald, V. *et al.* (2010) Rituximab pharmacokinetics during the management of acute idiopathic thrombotic thrombocytopenic purpura. <u>J Thromb Haemost. 8: 1201-8.</u>
 Kagan, L. *et al.* (2012) Subcutaneous Absorption of Monoclonal Antibodies: Role of Dose, Site of Injection, and Injection Volume on Rituximab Pharmacokinetics in Rats. <u>Pharm Res. 29: 490-499</u>

11. Kagan, L. and Mager, D.E. (2013) Mechanisms of subcutaneous absorption of rituximab in rats. <u>Drug Metab Dispos. 41: 248-55.</u>

12. Liu, X.F. *et al.* (2012) Validation of a Gyrolab<sup>™</sup> assay for quantification of rituximab in human serum. J Pharmacol Toxicol Methods. 65: 107-14.

13. Kagan, L. *et al.* (2014) Interspecies pharmacokinetic modeling of subcutaneous absorption of rituximab in mice and rats. <u>Pharm Res. 31: 3265-73.</u>

Blasco, H. *et al.* (2009) Pharmacokinetics of rituximab associated with CHOP chemotherapy in B-cell non-Hodgkin lymphoma. <u>Fundam Clin Pharmacol. 23: 601-8.</u>
 Illidge, T.M. *et al.* (2016) Short duration immunochemotherapy followed by radioimmunotherapy consolidation is effective and well tolerated in relapsed follicular lymphoma: 5-year results from a UK National Cancer Research Institute Lymphoma Group study. <u>Br J Haematol. 173 (2): 274-82.</u>

16. Vacher, P. *et al.* (2015) Localized Store-Operated Calcium Influx Represses CD95-Dependent Apoptotic Effects of Rituximab in Non-Hodgkin B Lymphomas. J Immunol. pii: 1402942.

17. Komori, M. *et al.* (2016) Insufficient disease inhibition by intrathecal rituximab in progressive multiple sclerosis. <u>Ann Clin Transl Neurol. 3 (3): 166-79.</u>

18. Kashiwagi, N. *et al.* (2017) Method for measuring anti-drug antibody <u>US Patent</u> <u>Application US20170315118A1</u>

19. Lioger, B. *et al.* (2017) Antigenic burden and serum IgG concentrations influence rituximab pharmacokinetics in rheumatoid arthritis patients. <u>Br J Clin Pharmacol. 83 (8):</u> 1773-81.

20. Zhang, Y. *et al.* (2013) Stability of stock and diluted rituximab. <u>Am J Health Syst</u> <u>Pharm. 70 (5): 436-8.</u>

21. Desbois, A.C. *et al.* (2020) Rituximab-associated Vasculitis Flare: Incidence, Predictors, and Outcome. <u>J Rheumatol. 47 (6): 896-902.</u>

StorageThis product is shipped at ambient temperature. It is recommended to aliquot and store at<br/>-20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for<br/>short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Acknowledgements	Rituxan is a trademark of Biogen Idec/Genentech in the USA. MabThera is a trademark of Roche in Europe.
Health And Safety Information	Material Safety Datasheet documentation #10057 available at: https://www.bio-rad-antibodies.com/SDS/MCA2260
Regulatory	For research purposes only

## Related Products

### **Recommended Secondary Antibodies**

Rabbit Anti Rat IgG (STAR20...) <u>RPE</u> Rabbit Anti Rat IgG (STAR17...) <u>FITC</u> Goat Anti Rat IgG (STAR69...) <u>FITC</u> Goat Anti Rat IgG (STAR73...) <u>RPE</u> **Recommended Useful Reagents** 

HUMAN ANTI RITUXIMAB (HCA061) HUMAN ANTI RITUXIMAB (HCA186)

### Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M443063:250619'

#### Printed on 19 Jun 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint