

## Datasheet: MCA2274PE

<b>Description:</b>	RAT ANTI MOUSE CD150:RPE
<b>Specificity:</b>	CD150
<b>Other names:</b>	SLAM
<b>Format:</b>	RPE
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	9D1
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	100 TESTS

### Product Details

#### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Mouse						
<b>Product Form</b>	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized						
<b>Reconstitution</b>	Reconstitute with 1 ml distilled water						
<b>Max Ex/Em</b>	<table border="1"> <thead> <tr> <th>Fluorophore</th> <th>Excitation Max (nm)</th> <th>Emission Max (nm)</th> </tr> </thead> <tbody> <tr> <td>RPE 488nm laser</td> <td>496</td> <td>578</td> </tr> </tbody> </table>	Fluorophore	Excitation Max (nm)	Emission Max (nm)	RPE 488nm laser	496	578
Fluorophore	Excitation Max (nm)	Emission Max (nm)					
RPE 488nm laser	496	578					
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant						
<b>Buffer Solution</b>	Phosphate buffered saline						
<b>Preservative</b>	0.09% Sodium Azide						
<b>Stabilisers</b>	1% Bovine Serum Albumin 5% Sucrose						

<b>Immunogen</b>	CHO cells stably transfected with mouse CD150
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">Q9QUM4</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">27218</a> Slamf1    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	Slam
<b>RRID</b>	AB_566533
<b>Fusion Partners</b>	Spleen cells from immunised rats were fused with cells of a myeloma cell line.
<b>Specificity</b>	<p><b>Rat anti Mouse CD150 antibody, clone 9D1</b> recognizes murine CD150, also known as signaling lymphocyte activation molecule (SLAM) . Murine CD150 is a 75 kDa cell surface glycoprotein that is expressed on T-cells and B-cells, and is upregulated on activated T-cells and stimulated macrophages. The extracellular domain of CD150 is the receptor for the measles virus (<a href="#">Erlenhoefer et al. 2001</a>) and acts as a co-activator on T-cells and B-cells.</p> <p>Rat anti Mouse CD150 antibody, clone 9D1 has functional activity (<a href="#">Howie et al. 2002</a>).</p>
<b>Flow Cytometry</b>	<p>Use 10ul of the suggested working dilution to label <math>10^6</math> cells in 100ul. &lt;</p> <p>The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity fc receptors. This may be reduced by using SeroBlock FcR (<a href="#">BUF041A/B</a>).</p>
<b>References</b>	<ol style="list-style-type: none"> <li>Howie, D. <i>et al.</i> (2002) The role of SAP in murine CD150 (SLAM)-mediated T-cell proliferation and interferon gamma production. <a href="#">Blood. 100 (8): 2899-907.</a></li> <li>Despots, C. <i>et al.</i> (2006) SHIP deficiency enhances HSC proliferation and survival but compromises homing and repopulation. <a href="#">Blood. 107: 4338-45.</a></li> <li>Almolda, B. <i>et al.</i> (2015) Alterations in microglial phenotype and hippocampal neuronal function in transgenic mice with astrocyte-targeted production of interleukin-10. <a href="#">Brain Behav Immun. 45: 80-97.</a></li> <li>Villacampa, N. <i>et al.</i> (2015) Astrocyte-targeted production of IL-10 induces changes in microglial reactivity and reduces motor neuron death after facial nerve axotomy. <a href="#">Glia. 63 (7): 1166-84.</a></li> </ol>
<b>Storage</b>	<p>Prior to reconstitution store at +4°C. Following reconstitution store at +4°C.</p> <p>DO NOT FREEZE.</p> <p>This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.</p>

**Guarantee** 12 months from date of despatch

---

**Health And Safety Information** Material Safety Datasheet documentation #20487 available at:  
20487: <https://www.bio-rad-antibodies.com/uploads/MSDS/20487.pdf>

---

**Regulatory** For research purposes only

---

## Related Products

### Recommended Useful Reagents

[MOUSE SEROBLOCK FcR \(BUF041A\)](#)

[MOUSE SEROBLOCK FcR \(BUF041B\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M375450:210104'

**Printed on 07 May 2021**

---

© 2021 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)