

Datasheet: MCA2277 BATCH NUMBER 163431

Description:	escription: MOUSE ANTI DENGUE VIRUS		
Specificity:	DENGUE VIRUS		
Format:	Purified		
Product Type:	Monoclonal Antibody		
Clone:	Dengue 1-11(3)		
Isotype:	lgG2a		
Quantity:	0.25 mg		

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				
Immunohistology - Frozen			•	
Immunohistology - Paraffin (1)				
ELISA	•			
Immunoprecipitation			•	
Western Blotting				
Immunofluorescence	-			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1)This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose.

Target Species	Viral	
Product Form	Purified IgG - liquid	
Preparation	Purified IgG prepared by affinity chromatography on Protein A supernatant	A from tissue culture
Buffer Solution	Phosphate buffered saline	

<0.1% Sodium Azide (NaN ₃)
IgG concentration 1.0mg/ml
AB_609623
Mouse anti-Dengue virus antibody, clone Dengue 1-11 (3) recognizes all four dengue virus serotypes (DEN-1, DEN-2, DEN-3 and DEN-4) of the genus <i>Flavivirus</i> .
Mouse anti-Dengue virus antibody, clone Dengue 1-11 (3) binds strongly to subtypes 1 and 2. It recognizes subtype 3 more weakly. Among the four subtypes, subtype 4 is recognized the least strongly and some customers have reported no recognition of subtype 4.
The dengue virus is responsible for the tropical and sub-tropical diseases, dengue (DF) and dengue haemorrhagic fever (DHF), transmitted to individuals through the bite of the <i>Aedes</i> mosquito. The global distribution of dengue is expanding and comparable to that of malaria, with symptoms ranging from a mild viral syndrome of high fever, rash, headache, fatigue and muscle and joint pain, to the more severe and sometimes fatal DHF. Each dengue virus serotype is antigenically distinct, such that infection only provides an individual with immunity to the causative serotype, with the possibility of further infection by a different serotype or even multiple serotypes.
This antibody detects an anti-E/envelope specific band of 61kDa under reducing conditions in Western blotting. A weak secondary band of 80kDa may also be apparent.
 Poggianella, M. et al. (2015) Dengue E Protein Domain III-Based DNA Immunisation Induces Strong Antibody Responses to All Four Viral Serotypes. PLoS Negl Trop Dis. 9 (7): e0003947. Slon Campos, J.L. (2017) Evaluation of a Tetravalent DNA Vaccine against Dengue: Integrating Biochemical Studies on Dengue Virus Envelope Protein to a Domain-Based Antigen Design. PhD thesis The Open University. Nguyen, N.L. et al. (2015) Expression and characterization of an M cell-specific ligand-fused dengue virus tetravalent epitope using Saccharomyces cerevisiae. J Biosci Bioeng. 119 (1): 19-27. Kim, T.G. et al. (2010) Cholera toxin B subunit-domain III of dengue virus envelope glycoprotein E fusion protein production in transgenic plants. Protein Expr Purif. 74:236-41. Milic, N.L. et el. (2015) Sequence analysis and characterisation of virally induced viperin in the saltwater crocodile (Crocodylus porosus). Dev Comp Immunol. 51 (1): 108-15. Kim, M.Y. et al. (2018) Plant-expressed Fc-fusion protein tetravalent dengue vaccine

with inherent adjuvant properties. Plant Biotechnol J. 16 (7): 1283-94.

infection Plant Cell, Tissue and Organ Culture (PCTOC). 137 (1): 1-10.

7. Kim, B. & Kim, M. (2019) Evaluation of the oral immunogenicity of M cell-targeted tetravalent EDIII antigen for development of plant-based edible vaccine against dengue

Further Reading

- 1. Gubler, D.J. (1998) Dengue and dengue hemorrhagic fever. Clin Microbiol Rev. 11 (3):
- 2. Loroño-pino, M.A. *et al.* (1999) Common occurrence of concurrent infections by multiple dengue virus serotypes. Am J Trop Med Hyg. 61 (5): 725-30.
- 3. PhilipSamuel, P. & Tyagi, B.K. (2006) Diagnostic methods for detection & isolation of dengue viruses from vector mosquitoes. <u>Indian J Med Res. 123 (5): 615-28.</u>

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA2277 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)

Goat Anti Mouse IgG IgA IgM (STAR87...)

RPE

Goat Anti Mouse IgG (STAR76...)

RPE

Rabbit Anti Mouse IgG (STAR13...)

HRP

Goat Anti Mouse IgG (STAR70...)

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®550,

DyLight®650, DyLight®680, DyLight®800,

FITC, HRP

Rabbit Anti Mouse IgG (STAR9...) FITC

Goat Anti Mouse IgG (STAR77...) HRP

Goat Anti Mouse IgG (Fc) (STAR120...) FITC, HRP

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739

Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

-rad.com Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M389413:210806'

Printed on 05 Feb 2024