

Datasheet: MCA2277

**BATCH NUMBER 167589**

<b>Description:</b>	MOUSE ANTI DENGUE VIRUS
<b>Specificity:</b>	DENGUE VIRUS
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	Dengue 1-11(3)
<b>Isotype:</b>	IgG2a
<b>Quantity:</b>	0.25 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin (1)	▪			
ELISA	▪			
Immunoprecipitation			▪	
Western Blotting	▪			
Immunofluorescence	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

**(1)This product requires antigen retrieval using heat treatment prior to staining of paraffin sections.Sodium citrate buffer pH 6.0 is recommended for this purpose.**

<b>Target Species</b>	Viral
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline

<b>Preservative Stabilisers</b>	<0.1% sodium azide (NaN <sub>3</sub> )
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0mg/ml
<b>RRID</b>	AB_609623
<b>Specificity</b>	<p><b>Mouse anti-Dengue virus antibody, clone Dengue 1-11 (3)</b> recognizes all four dengue virus serotypes (DEN-1, DEN-2, DEN-3 and DEN-4) of the genus <i>Flavivirus</i>.</p> <p>Mouse anti-Dengue virus antibody, clone Dengue 1-11 (3) binds strongly to subtypes 1 and 2. It recognizes subtype 3 more weakly. Among the four subtypes, subtype 4 is recognized the least strongly and some customers have reported no recognition of subtype 4.</p> <p>The dengue virus is responsible for the tropical and sub-tropical diseases, dengue (DF) and dengue haemorrhagic fever (DHF), transmitted to individuals through the bite of the <i>Aedes</i> mosquito. The global distribution of dengue is expanding and comparable to that of malaria, with symptoms ranging from a mild viral syndrome of high fever, rash, headache, fatigue and muscle and joint pain, to the more severe and sometimes fatal DHF. Each dengue virus serotype is antigenically distinct, such that infection only provides an individual with immunity to the causative serotype, with the possibility of further infection by a different serotype or even multiple serotypes.</p>
<b>Western Blotting</b>	This antibody detects an anti-E/envelope specific band of ~61 kDa under reducing conditions. A weak secondary band of ~80 kDa may also be apparent.
<b>References</b>	<ol style="list-style-type: none"> <li>1. Kim, T.G. <i>et al.</i> (2010) Cholera toxin B subunit-domain III of dengue virus envelope glycoprotein E fusion protein production in transgenic plants. <a href="#">Protein Expr Purif. 74:236-41.</a></li> <li>2. Nguyen, N.L. <i>et al.</i> (2015) Expression and characterization of an M cell-specific ligand-fused dengue virus tetravalent epitope using <i>Saccharomyces cerevisiae</i>. <a href="#">J Biosci Bioeng. 119 (1): 19-27.</a></li> <li>3. Poggianella, M. <i>et al.</i> (2015) Dengue E Protein Domain III-Based DNA Immunisation Induces Strong Antibody Responses to All Four Viral Serotypes. <a href="#">PLoS Negl Trop Dis. 9 (7): e0003947.</a></li> <li>4. Milic, N.L. <i>et al.</i> (2015) Sequence analysis and characterisation of virally induced viperin in the saltwater crocodile (<i>Crocodylus porosus</i>). <a href="#">Dev Comp Immunol. 51 (1): 108-15.</a></li> <li>5. Slon Campos, J.L. (2017) Evaluation of a Tetravalent DNA Vaccine against Dengue: Integrating Biochemical Studies on Dengue Virus Envelope Protein to a Domain-Based Antigen Design. <a href="#">PhD thesis The Open University.</a></li> <li>6. Kim, M.Y. <i>et al.</i> (2018) Plant-expressed Fc-fusion protein tetravalent dengue vaccine with inherent adjuvant properties. <a href="#">Plant Biotechnol J. 16 (7): 1283-94.</a></li> <li>7. Kim, B. &amp; Kim, M. (2019) Evaluation of the oral immunogenicity of M cell-targeted tetravalent EDIII antigen for development of plant-based edible vaccine against dengue infection <a href="#">Plant Cell, Tissue and Organ Culture (PCTOC). 137 (1): 1-10.</a></li> </ol>

8. Kim, M.Y. *et al.* (2024) Marked enhancement of the immunogenicity of plant-expressed IgG-Fc fusion proteins by inclusion of cholera toxin non-toxic B subunit within the single polypeptide. [Plant Biotechnol J. Jan 01 \[Epub ahead of print\]](#).

- Further Reading**
1. Gubler, D.J. (1998) Dengue and dengue hemorrhagic fever. [Clin Microbiol Rev. 11 \(3\): 480-96.](#)
  2. Loroño-pino, M.A. *et al.* (1999) Common occurrence of concurrent infections by multiple dengue virus serotypes. [Am J Trop Med Hyg. 61 \(5\): 725-30.](#)
  3. PhilipSamuel, P. & Tyagi, B.K. (2006) Diagnostic methods for detection & isolation of dengue viruses from vector mosquitoes. [Indian J Med Res. 123 \(5\): 615-28.](#)

**Storage** This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

**Guarantee** 12 months from date of despatch

**Health And Safety Information** Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA2277>  
10040

**Regulatory** For research purposes only

## Related Products

### Recommended Secondary Antibodies

- Rabbit Anti Mouse IgG (STAR12...) [RPE](#)  
Goat Anti Mouse IgG IgA IgM (STAR87...) [HRP](#)  
Goat Anti Mouse IgG (STAR76...) [RPE](#)  
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)  
Goat Anti Mouse IgG (STAR70...) [FITC](#)  
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®550](#),  
[DyLight®650](#), [DyLight®680](#), [DyLight®800](#),  
[FITC](#), [HRP](#)  
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)  
Goat Anti Mouse IgG (STAR77...) [HRP](#)  
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)

**North & South America** Tel: +1 800 265 7376  
Fax: +1 919 878 3751  
Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide** Tel: +44 (0)1865 852 700  
Fax: +44 (0)1865 852 739  
Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe** Tel: +49 (0) 89 8090 95 21  
Fax: +49 (0) 89 8090 95 50  
Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M418780:230427'

Printed on 05 Feb 2024

