

Datasheet: MCA2281A647

BATCH NUMBER 166583

Description:	RAT ANTI MOUSE CD200R:Alexa Fluor® 647
Specificity:	CD200R
Other names:	OX2R
Format:	ALEXA FLUOR® 647
Product Type:	Monoclonal Antibody
Clone:	OX-110
Isotype:	IgG2a
Quantity:	100 TESTS/1ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse						
Product Form	Purified IgG conjugated to Alexa Fluor® 647 - liquid						
Max Ex/Em	<table border="1"> <thead> <tr> <th>Fluorophore</th> <th>Excitation Max (nm)</th> <th>Emission Max (nm)</th> </tr> </thead> <tbody> <tr> <td>Alexa Fluor®647</td> <td>650</td> <td>665</td> </tr> </tbody> </table>	Fluorophore	Excitation Max (nm)	Emission Max (nm)	Alexa Fluor®647	650	665
Fluorophore	Excitation Max (nm)	Emission Max (nm)					
Alexa Fluor®647	650	665					
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant						
Buffer Solution	Phosphate buffered saline						
Preservative	0.09% sodium azide (NaN ₃)						
Stabilisers	1% bovine serum albumin						
Approx. Protein	IgG concentration 0.05 mg/ml						

Concentrations

Immunogen Fusion protein mCD200RCD4d3+4

External Database Links

UniProt:

[Q9ES57](#) [Related reagents](#)

Entrez Gene:

[57781](#) Cd200r1 [Related reagents](#)

Synonyms Mox2r, Ox2r

RRID AB_566615

Specificity

Rat anti Mouse CD200R antibody, clone OX-110 recognizes mouse CD200R, a cell surface glycoprotein (also known as OX2R. Mouse CD200R is a 326 amino acid, ~48 kDa single pass type I transmembrane glycoprotein, expressed primarily by peripheral blood monocytes and neutrophils but also by other leucocytes including T-lymphocytes and mast cells ([Wright et al.2003](#)). CD200-CD200R interaction may be involved in the control of myeloid cellular function ([Minas and Liversidge 2006](#)).

Rat anti Mouse CD200R antibody, clone OX-110 has been used successfully for the immunohistochemical detection of CD200R expressing cells on acetone fixed cryosections of murine synovial tissue ([Simelyte et al. 2008](#)).

Flow Cytometry

Use 10µl of the suggested working dilution to label 10⁶ cells in 100µl.

The Fc region of monoclonal antibodies may bind to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR ([BUF041A/BUF041B](#)).

References

1. Wright, G.J. *et al.* (2003) Characterization of the CD200 receptor family in mice and humans and their interactions with CD200. [J Immunol. 171 \(6\): 3034-46.](#)
2. Zhao, J. *et al.* (2009) Evasion by stealth: inefficient immune activation underlies poor T cell response and severe disease in SARS-CoV-infected mice. [PLoS Pathog. 5 \(10\): e1000636.](#)
3. Björck, P. *et al.* (2011) Plasmacytoid Dendritic Cell Dichotomy: Identification of IFN- α Producing Cells as a Phenotypically and Functionally Distinct Subset. [J Immunol. 186: 1477-85.](#)
4. Dellacasa-Lindberg, I. *et al.* (2011) Migratory Activation of Primary Cortical Microglia upon Infection with *Toxoplasma gondii*. [Infect Immun. 79: 3046-52.](#)
5. Krejsek, J. *et al.* (2010) Expression of CD200/CD200R regulatory molecules on granulocytes and monocytes is modulated by cardiac surgical operation. [Perfusion 25: 389-97.](#)
6. Rijkers, E.S. *et al.* (2008) The inhibitory CD200R is differentially expressed on human and mouse T and B lymphocytes. [Mol Immunol. 45: 1126-35.](#)
7. Costello, D.A. *et al.* (2011) Long term potentiation is impaired in membrane glycoprotein CD200-deficient mice: a role for Toll-like receptor activation. [J Biol Chem. 286 \(40\): 34722-32.](#)

8. Simelyte, E. *et al.* (2008) CD200-Fc, a novel antiarthritic biologic agent that targets proinflammatory cytokine expression in the joints of mice with collagen-induced arthritis. [Arthritis Rheum. 58: 1038-43.](#)
9. Whitmore, L.C. *et al.* (2014) NOX2 protects against progressive lung injury and multiple organ dysfunction syndrome. [Am J Physiol Lung Cell Mol Physiol. 307: L71-82.](#)
10. Chitnis, T. *et al.* (2007) Elevated neuronal expression of CD200 protects Wlds mice from inflammation-mediated neurodegeneration. [Am J Pathol. 170: 1695-712.](#)
11. Larson, D. *et al.* (2012) Chronic helminth infection reduces basophil responsiveness in an IL-10-dependent manner. [J Immunol. 188: 4188-99.](#)
12. Torrero, M.N. *et al.* (2009) CD200R surface expression as a marker of murine basophil activation. [Clin Exp Allergy. 39 \(3\): 361-9.](#)
13. Hernangómez M *et al.* (2012) CD200-CD200R1 interaction contributes to neuroprotective effects of anandamide on experimentally induced inflammation. [Glia 60 \(9\): 1437-50.](#)
14. Bain, C.C. & Mowat, A.M. (2012) CD200 receptor and macrophage function in the intestine. [Immunobiology 217 \(6\): 643-51.](#)
15. Seeds, R.E. *et al.* (2011) The role of myeloid receptors on murine plasmacytoid dendritic cells in induction of type I interferon. [Int Immunopharmacol. 11 \(7\): 794-801.](#)
16. Akkaya, M. *et al.* (2013) Dissection of agonistic and blocking effects of CD200 receptor antibodies. [PLoS One 8 \(5\): e63325.](#)
17. Cassard L *et al.* (2012) Fcγ receptors inhibit mouse and human basophil activation. [J Immunol. 189 \(6\): 2995-3006.](#)
18. Liu, J.Q. *et al.* (2016) A Critical Role for CD200R Signaling in Limiting the Growth and Metastasis of CD200+ Melanoma. [J Immunol. Jul 6. pii: 1600052. \[Epub ahead of print\]](#)
19. Toya, E.E. & Ohba, M. (2015) Improved Long-term Culture of Epidermal Stem Cells Utilizing CD200R-expressing Feeder Cells [Showa Uni J Med Sci 27 \(2\) 83-91.](#)
20. Lago, N. *et al.* (2018) CD200 modulates spinal cord injury neuroinflammation and outcome through CD200R1. [Brain Behav Immun. 73: 416-26.](#)
21. Qian, H. *et al.* (2023) Activation of the CD200/CD200R1 axis attenuates neuroinflammation and improves postoperative cognitive dysfunction via the PI3K/Akt/NF-κB signaling pathway in aged mice. [Inflamm Res. 72 \(12\): 2127-44.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee 12 months from date of despatch

Acknowledgements This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchase product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing

or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or outlicensing@thermofisher.com

Health And Safety Information

Material Safety Datasheet documentation #10041 available at:
<https://www.bio-rad-antibodies.com/SDS/MCA2281A647>
10041

Regulatory

For research purposes only

Related Products

Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL:Alexa Fluor® 647 \(MCA1212A647\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M413940:221129'

Printed on 23 May 2025

© 2025 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)