

Datasheet: MCA2281GA

BATCH NUMBER 158699

Description:	RAT ANTI MOUSE CD200R
Specificity:	CD200R
Other names:	OX2R
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	OX-110
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/25 - 1/50
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide

Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Fusion protein mCD200RCD4d3+4
External Database Links	<p>UniProt: Q9ES57 Related reagents</p> <p>Entrez Gene: 57781 Cd200r1 Related reagents</p>
Synonyms	Mox2r, Ox2r
RRID	AB_566623
Specificity	<p>Rat anti Mouse CD200R antibody, clone OX-110 recognizes mouse CD200R, a cell surface glycoprotein (also known as OX2R. Mouse CD200R is a 326 amino acid, ~48 kDa single pass type I transmembrane glycoprotein, expressed primarily by peripheral blood monocytes and neutrophils but also by other leucocytes including T-lymphocytes and mast cells (Wright et al.2003). CD200-CD200R interaction may be involved in the control of myeloid cellular function (Minas and Liversidge 2006).</p> <p>Rat anti Mouse CD200R antibody, clone OX-110 has been used successfully for the immunohistochemical detection of CD200R expressing cells on acetone fixed cryosections of murine synovial tissue (Simelyte et al. 2008).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul
References	<ol style="list-style-type: none"> 1. Wright, G.J. <i>et al.</i> (2003) Characterization of the CD200 receptor family in mice and humans and their interactions with CD200. J Immunol. 171 (6): 3034-46. 2. Zhao, J. <i>et al.</i> (2009) Evasion by stealth: inefficient immune activation underlies poor T cell response and severe disease in SARS-CoV-infected mice. PLoS Pathog. 5 (10): e1000636. 3. Björck, P. <i>et al.</i> (2011) Plasmacytoid Dendritic Cell Dichotomy: Identification of IFN-α Producing Cells as a Phenotypically and Functionally Distinct Subset. J Immunol. 186: 1477-85. 4. Dellacasa-Lindberg, I. <i>et al.</i> (2011) Migratory Activation of Primary Cortical Microglia upon Infection with <i>Toxoplasma gondii</i>. Infect Immun. 79: 3046-52. 5. Krejsek, J. <i>et al.</i> (2010) Expression of CD200/CD200R regulatory molecules on granulocytes and monocytes is modulated by cardiac surgical operation. Perfusion 25: 389-97. 6. Rijkers, E.S. <i>et al.</i> (2008) The inhibitory CD200R is differentially expressed on human and mouse T and B lymphocytes. Mol Immunol. 45: 1126-35. 7. Costello, D.A. <i>et al.</i> (2011) Long term potentiation is impaired in membrane glycoprotein CD200-deficient mice: a role for Toll-like receptor activation. J Biol Chem. 286 (40):

[34722-32.](#)

8. Simelyte, E. *et al.* (2008) CD200-Fc, a novel antiarthritic biologic agent that targets proinflammatory cytokine expression in the joints of mice with collagen-induced arthritis. [Arthritis Rheum. 58: 1038-43.](#)
9. Whitmore, L.C. *et al.* (2014) NOX2 protects against progressive lung injury and multiple organ dysfunction syndrome. [Am J Physiol Lung Cell Mol Physiol. 307: L71-82.](#)
10. Chitnis, T. *et al.* (2007) Elevated neuronal expression of CD200 protects Wilds mice from inflammation-mediated neurodegeneration. [Am J Pathol. 170: 1695-712.](#)
11. Larson, D. *et al.* (2012) Chronic helminth infection reduces basophil responsiveness in an IL-10-dependent manner. [J Immunol. 188: 4188-99.](#)
12. Torrero, M.N. *et al.* (2009) CD200R surface expression as a marker of murine basophil activation. [Clin Exp Allergy. 39 \(3\): 361-9.](#)
13. Hernangómez M *et al.* (2012) CD200-CD200R1 interaction contributes to neuroprotective effects of anandamide on experimentally induced inflammation. [Glia 60 \(9\): 1437-50.](#)
14. Bain, C.C. & Mowat, A.M. (2012) CD200 receptor and macrophage function in the intestine. [Immunobiology 217 \(6\): 643-51.](#)
15. Seeds, R.E. *et al.* (2011) The role of myeloid receptors on murine plasmacytoid dendritic cells in induction of type I interferon. [Int Immunopharmacol. 11 \(7\): 794-801.](#)
16. Akkaya, M. *et al.* (2013) Dissection of agonistic and blocking effects of CD200 receptor antibodies. [PLoS One 8 \(5\): e63325.](#)
17. Cassard L *et al.* (2012) Fcγ receptors inhibit mouse and human basophil activation. [J Immunol. 189 \(6\): 2995-3006.](#)
18. Liu, J.Q. *et al.* (2016) A Critical Role for CD200R Signaling in Limiting the Growth and Metastasis of CD200+ Melanoma. [J Immunol. Jul 6. pii: 1600052. \[Epub ahead of print\]](#)
19. Toya, E.E. & Ohba, M. (2015) Improved Long-term Culture of Epidermal Stem Cells Utilizing CD200R-expressing Feeder Cells [Showa Uni J Med Sci 27 \(2\) 83-91.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA2281GA>
10040

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16...)

[DyLight®800](#)

Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (STAR72...)	HRP
Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR73...)	RPE
Rabbit Anti Rat IgG (STAR21...)	HRP
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight®550 , DyLight®650 , DyLight®800
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin

Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL \(MCA1212\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M383675:210513'

Printed on 21 Feb 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)