

Datasheet: MCA2282A647

BATCH NUMBER 166481

Description:	MOUSE ANTI HUMAN CD200R:Alexa Fluor® 647
Specificity:	CD200R
Other names:	OX2R
Format:	ALEXA FLUOR® 647
Product Type:	Monoclonal Antibody
Clone:	OX108
Isotype:	IgG1
Quantity:	100 TESTS/1ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG conjugated to Alexa Fluor® 647 - liquid

Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	Alexa Fluor®647	650	665

Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
--------------------	---

Buffer Solution	Phosphate buffered saline
------------------------	---------------------------

Preservative	0.09% sodium azide (NaN ₃)
Stabilisers	1% bovine serum albumin

Approx. Protein	IgG concentration 0.05 mg/ml
------------------------	------------------------------

Concentrations

Immunogen Fusion protein hCD200RCD4d3+4.

External Database Links

UniProt:

[Q8TD46](#) [Related reagents](#)

Entrez Gene:

[131450](#) CD200R1 [Related reagents](#)

Synonyms CD200R, CRTR2, MOX2R, OX2R

RRID AB_566616

Specificity

Mouse anti Human CD200R antibody, clone OX108 recognizes human CD200R, a cell surface glycoprotein (also known as OX2R). In humans CD200R is expressed primarily by peripheral blood monocytes and neutrophils but also by other leucocytes including T-lymphocytes and mast cells, CD200-CD200R interaction may be involved in the control of myeloid cellular function ([Wright *et al.* 2003](#)).

Levels of expression on resting peripheral blood cells are relatively low.

Flow Cytometry

Use 10µl of the suggested working dilution to label 10⁶ cells in 100µl

References

1. Wright, G.J. *et al.* (2003) Characterization of the CD200 receptor family in mice and humans and their interactions with CD200. [J Immunol. 171 \(6\): 3034-46.](#)
2. Koning, N. *et al.* (2009) Distribution of the immune inhibitory molecules CD200 and CD200R in the normal central nervous system and multiple sclerosis lesions suggests neuron-glia and glia-glia interactions. [J Neuropathol Exp Neurol. 68: 159-67.](#)
3. Melief, J. *et al.* (2012) Phenotyping primary human microglia: tight regulation of LPS responsiveness. [Glia. 60 \(10\): 1506-17.](#)
4. Ambarus CA *et al.* (2012) Soluble immune complexes shift the TLR-induced cytokine production of distinct polarized human macrophage subsets towards IL-10. [PLoS One. 7 \(4\): e35994.](#)
5. Ambarus, C.A. *et al.* (2012) Intimal lining layer macrophages but not synovial sublining macrophages display an IL-10 polarized-like phenotype in chronic synovitis. [Arthritis Res Ther. 14 \(2\): R74.](#)
6. Vicetti, R.M. *et al.* (2013) Human female genital tract infection by the obligate intracellular bacterium *Chlamydia trachomatis* elicits robust Type 2 immunity. [PLoS One. 8 \(3\): e58565.](#)
7. Melief, J. *et al.* (2013) Microglia in normal appearing white matter of multiple sclerosis are alerted but immunosuppressed. [Glia. 61 \(11\): 1848-61.](#)
8. Vicetti Miguel, R.D. *et al.* (2013) Human female genital tract infection by the obligate intracellular bacterium *Chlamydia trachomatis* elicits robust Type 2 immunity. [PLoS One. 8 \(3\): e58565.](#)
9. Memarian, A. *et al.* (2013) Upregulation of CD200 is associated with Foxp3+ regulatory

T cell expansion and disease progression in acute myeloid leukemia. [Tumour Biol. 34 \(1\): 531-42.](#)

10. van de Garde, M.D. *et al.* (2014) Chronic exposure to glucocorticoids shapes gene expression and modulates innate and adaptive activation pathways in macrophages with distinct changes in leukocyte attraction. [J Immunol. 192 \(3\): 1196-208.](#)

11. Sousa, S. *et al.* (2015) Human breast cancer cells educate macrophages toward the M2 activation status. [Breast Cancer Res. 17 \(1\): 101.](#)

12. Holmannova, D. *et al.* (2015) Inhibitory CD200R and proapoptotic CD95/CD95L molecules on innate immunity cells are modulated by cardiac surgery. [Perfusion. 30 \(7\): 543-55.](#)

13. Wiedemann, G.M. *et al.* (2016) A novel TLR7 agonist reverses NK cell anergy and cures RMA-S lymphoma-bearing mice. [Oncoimmunology. 5 \(7\): e1189051.](#)

14. Björnfort Holmström, S. *et al.* (2017) Gingival Tissue Inflammation Promotes Increased Matrix Metalloproteinase-12 Production by CD200R^{low} Monocyte-Derived Cells in Periodontitis. [J Immunol. 199 \(12\): 4023-35.](#)

15. Pascoal, M.R. *et al.* (2022) CD200R1L is a functional evolutionary conserved activating receptor in human neutrophils. [J Leukoc Biol. 111 \(2\): 367-77.](#)

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee

12 months from date of despatch

Acknowledgements

This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchase product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or outlicensing@thermofisher.com

Health And Safety Information

Material Safety Datasheet documentation #10041 available at: <https://www.bio-rad-antibodies.com/SDS/MCA2282A647>
10041

Regulatory

For research purposes only

Related Products**Recommended Negative Controls**

[MOUSE IgG1 NEGATIVE CONTROL:Alexa Fluor® 647 \(MCA928A647\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M429713:240416'

Printed on 16 Apr 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)