

Datasheet: MCA2282B

Description:	MOUSE ANTI HUMAN CD200R:Biotin
Specificity:	CD200R
Other names:	OX2R
Format:	Biotin
Product Type:	Monoclonal Antibody
Clone:	OX108
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG conjugated to Biotin - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative	0.09% Sodium Azide
Stabilisers	1% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml
Immunogen	Fusion protein hCD200RCD4d3+4.
External Database Links	<p>UniProt: Q8TD46 Related reagents</p> <p>Entrez Gene: 131450 CD200R1 Related reagents</p>

Synonyms	CD200R, CRTR2, MOX2R, OX2R
RRID	AB_566618
Specificity	<p>Mouse anti Human CD200R antibody, clone OX108 recognizes human CD200R, a cell surface glycoprotein (also known as OX2R). In humans CD200R is expressed primarily by peripheral blood monocytes and neutrophils but also by other leucocytes including T-lymphocytes and mast cells, CD200-CD200R interaction may be involved in the control of myeloid cellular function (Wright et al. 2003).</p> <p>Levels of expression on resting peripheral blood cells are relatively low.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul
References	<ol style="list-style-type: none"> 1. Wright, G.J. <i>et al.</i> (2003) Characterization of the CD200 receptor family in mice and humans and their interactions with CD200. J Immunol. 171 (6): 3034-46. 2. Koning, N. <i>et al.</i> (2009) Distribution of the immune inhibitory molecules CD200 and CD200R in the normal central nervous system and multiple sclerosis lesions suggests neuron-glia and glia-glia interactions. J Neuropathol Exp Neurol. 68: 159-67. 3. Melief, J. <i>et al.</i> (2013) Microglia in normal appearing white matter of multiple sclerosis are alerted but immunosuppressed. Glia. 61 (11): 1848-61. 4. Ambarus, C.A. <i>et al.</i> (2012) Intimal lining layer macrophages but not synovial sublining macrophages display an IL-10 polarized-like phenotype in chronic synovitis. Arthritis Res Ther. 14 (2): R74. 5. Holmannova, D. <i>et al.</i> (2015) Inhibitory CD200R and proapoptotic CD95/CD95L molecules on innate immunity cells are modulated by cardiac surgery. Perfusion. 30 (7): 543-55. 6. Ambarus CA <i>et al.</i> (2012) Soluble immune complexes shift the TLR-induced cytokine production of distinct polarized human macrophage subsets towards IL-10. PLoS One. 7 (4): e35994. 7. Memarian, A. <i>et al.</i> (2013) Upregulation of CD200 is associated with Foxp3+ regulatory T cell expansion and disease progression in acute myeloid leukemia. Tumour Biol. 34 (1): 531-42. 8. Sousa, S. <i>et al.</i> (2015) Human breast cancer cells educate macrophages toward the M2 activation status. Breast Cancer Res. 17 (1): 101. 9. Vicetti Miguel, R.D. <i>et al.</i> (2013) Human female genital tract infection by the obligate intracellular bacterium <i>Chlamydia trachomatis</i> elicits robust Type 2 immunity. PLoS One. 8 (3): e58565. 10. van de Garde, M.D. <i>et al.</i> (2014) Chronic exposure to glucocorticoids shapes gene expression and modulates innate and adaptive activation pathways in macrophages with distinct changes in leukocyte attraction. J Immunol. 192 (3): 1196-208. 11. Melief, J. <i>et al.</i> (2012) Phenotyping primary human microglia: tight regulation of LPS responsiveness. Glia. 60 (10): 1506-17. 12. Georgoudaki, A.M. <i>et al.</i> (2016) Reprogramming Tumor-Associated Macrophages by Antibody Targeting Inhibits Cancer Progression and Metastasis. Cell Rep. 15 (9): 2000-11. 13. BjörnfortHolmström, S. <i>et al.</i> (2017) Gingival Tissue Inflammation Promotes Increased Matrix Metalloproteinase-12 Production by CD200R^{low} Monocyte-Derived Cells in Periodontitis. J Immunol. 199 (12): 4023-35. 14. Wiedemann, G.M. <i>et al.</i> (2016) A novel TLR7 agonist reverses NK cell anergy and cures RMA-S lymphoma-bearing mice. Oncoimmunology. 5 (7): e1189051.
Storage	<p>Store at +4°C or at -20°C if preferred.</p> <p>This product should be stored undiluted.</p> <p>Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this</p>

may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	18 months from date of despatch.
------------------	----------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M353734:190523'

Printed on 01 May 2020

© 2020 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)