

Datasheet: MCA2286

Description:	RAT ANTI MOUSE CD137L
Specificity:	CD137L
Other names:	4-1BBL, TNFSF9
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	AT113-2
Isotype:	IgG1
Quantity:	0.25 mg

Product Details

RRID AB_324889

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10 - 1/50
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species

Mouse

Product Form

Purified IgG - liquid

Preparation

Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant.

Buffer Solution

Phosphate buffered saline

Preservative Stabilisers

0.09% Sodium Azide

Carrier Free

Yes

Approx. Protein Concentrations

IgG concentration 1.0 mg/ml

Immunogen CD137L-Fc Fusion Protein.

External Database Links

UniProt:
[P41274](#) [Related reagents](#)

Entrez Gene:
[21950](#) Tnfsf9 [Related reagents](#)

Synonyms Cd137l, Cd157l, Ly63l

Specificity **Rat anti Mouse CD137L antibody, clone AT113-2** recognizes murine CD137L (4-1BBL), a 50 kDa type 2 transmembrane protein, which is a member of the TNF superfamily. In mice CD137L is expressed on activated T-cells, B-cells, macrophages and dendritic cells. Interaction of CD137L with its receptor CD137 (4-1BB) is reported to be involved in T and B cell growth and T cell effector function.

Flow Cytometry Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

References

1. Goodwin, R.G. *et al.* (1993) Molecular cloning of a ligand for the inducible T cell gene 4-1BB: a member of an emerging family of cytokines with homology to tumor necrosis factor. [Eur J Immunol. 23 \(10\): 2631-41.](#)
2. French, R.R. *et al.* (2007) Eradication of lymphoma by CD8 T cells following anti-CD40 monoclonal antibody therapy is critically dependent on CD27 costimulation. [Blood. 109 \(11\): 4810-5.](#)

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 18 months from date of despatch.

Health And Safety Information Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin
Goat Anti Rat IgG (STAR73...)	RPE
Rabbit Anti Rat IgG (STAR21...)	HRP
Goat Anti Rat IgG (STAR72...)	HRP
Rabbit Anti Rat IgG (STAR16...)	DyLight®800

Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...) [DyLight®800](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M341533:190109'

Printed on 20 May 2019

© 2019 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)