

Datasheet: MCA2289

BATCH NUMBER 166552

Description:	Description: RAT ANTI MOUSE DECTIN-	
Specificity:	DECTIN-1	
Other names:	CD369	
Format:	Purified	
Product Type:	Monoclonal Antibody	
Clone:	2A11	
Isotype:	lgG2b	
Quantity:	0.25 mg	

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				1/10 - 1/50
Immunohistology - Frozen (1)	•			
Immunohistology - Paraffin		•		
Immunohistology - Resin		•		
ELISA				
Immunoprecipitation	•			
Western Blotting			•	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) The epitope recognized by 2A11 is sensitive to H_20_2 treatment, therefore quenching with 0.3% H_20_2 should be performed after incubation with the primary antibody. The epitope recognized by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. Bio-Rad recommends the use of acetone fixation for frozen sections.

Target Species	Mouse
Product Form	Purified IgG - liquid

Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% sodium azide (NaN ₃)
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Dectin-1 transfected NIH3T3 cells and recombinant soluble Dectin-1.
External Database Links	UniProt: Q6QLQ4 Related reagents Entrez Gene: 56644 Clec7a Related reagents
Synonyms	Bgr, Clecsf12, Dectin1
RRID	AB_324904
Fusion Partners	Spleen cells from immunized Fischer rats were fused with cells of the rat Y3 myeloma cell line
Specificity	Rat anti Mouse dectin-1 antibody, clone 2A11 recognizes murine beta-glucan receptor, also known as dectin-1 or CD369. Dectin-1 is predominantly expressed by cells of the monocyte/macrophage and neutrophil lineages, but also at lower levels by dendritic cells and a subpopulation of T cells.
	As a major leucocyte receptor for beta-glucan this molecule may have a key role in the immunomodulatory effects of beta-glucans and in the host response to fungal pathogens. Dectin-1 may stimulate reactive oxygen production in macropahges via the protein tyrosine kinase known as Syk.
	Rat anti Mouse Dectin-1 antibody, clone 2A11 inhibits the binding of zymosan to macrophages via the beta-glucan receptor.
Flow Cytometry	Use 10µl of the suggested working dilution to label 10 ⁶ cells in 100µl
References	1. Taylor, P.R. <i>et al.</i> (2002) The beta-glucan receptor, dectin-1, is predominantly expressed on the surface of cells of the monocyte/macrophage and neutrophil lineages. <u>J Immunol.</u> 169 (7): 3876-82. 2. Reid, D.M. <i>et al.</i> (2004) Expression of the beta-glucan receptor, Dectin-1, on murine

leukocytes in situ correlates with its function in pathogen recognition and reveals potential

- roles in leukocyte interactions. J Leukoc Biol. 76 (1): 86-94.
- 3. Underhill, D.M. *et al.* (2005) Dectin-1 activates Syk tyrosine kinase in a dynamic subset of macrophages for reactive oxygen production. <u>Blood</u>. 106 (7): 2543-50.
- 4. Lefevre,L. *et al.* (2010) PPARc Ligands Switched High Fat Diet-Induced Macrophage M2b Polarization toward M2a Thereby Improving Intestinal *Candida* Elimination <u>PLoS</u> One. 5(9):e12828.
- 5. Fei, M. *et al.* (2011) TNF-alpha from inflammatory dendritic cells (DCs) regulates lung IL-17A/IL-5 levels and neutrophilia versus eosinophilia during persistent fungal infection. Proc Natl Acad Sci U S A. 108 (13): 5360-5.
- 6. Gazi, U. *et al.* (2011) Fungal Recognition Enhances Mannose Receptor Shedding through Dectin-1 Engagement. J Biol Chem. 286: 7822-9.
- 7. McDonald, J.U. *et al.* (2011) *In vivo* functional analysis and genetic modification of *in vitro*-derived mouse neutrophils. <u>FASEB J. 25 (6)</u>: 1972-82.
- 8. Dewals, B.G. *et al.* (2010) IL-4Ralpha-independent expression of mannose receptor and Ym1 by macrophages depends on their IL-10 responsiveness. <u>PLoS Negl Trop Dis. 4:</u> e689.
- 9. Galès, A. *et al.* (2010) PPARgamma controls dectin-1 expression required for host antifungal defense against *Candida albicans*. <u>PLoS Pathog. 6: e1000714</u>.
- 10. Goodridge, H.S. *et al.* (2009) Differential use of CARD9 by dectin-1 in macrophages and dendritic cells. <u>J Immunol</u>. 182: 1146-54.
- 11. Coates, P.J. *et al.* (2008) Indirect macrophage responses to ionizing radiation: implications for genotype-dependent bystander signaling. <u>Cancer Res. 68: 450-6.</u>
- 12. Dioszeghy, V. *et al.* (2008) 12/15-Lipoxygenase regulates the inflammatory response to bacterial products *in vivo*. <u>J Immunol. 181: 6514-24.</u>
- 13. Hohl, T.M. (2008) Caspofungin modulates inflammatory responses to *Aspergillus fumigatus* through stage-specific effects on fungal beta-glucan exposure. <u>J Infect Dis. 198:</u> 176-85.
- 14. Palma, A.S. *et al.* (2006) Ligands for the beta-glucan receptor, Dectin-1, assigned using "designer" microarrays of oligosaccharide probes (neoglycolipids) generated from glucan polysaccharides. <u>J Biol Chem. 281: 5771-9.</u>
- 15. Sindrilaru, A. *et al.* (2011) An unrestrained proinflammatory M1 macrophage population induced by iron impairs wound healing in humans and mice. <u>J Clin Invest. 121:</u> 985-97.
- 16. Anandasabapathy, N. *et al.* (2011) Flt3L controls the development of radiosensitive dendritic cells in the meninges and choroid plexus of the steady-state mouse brain. <u>J Exp</u> Med. 208 (8): 1695-705.
- 17. Takahara, K. *et al.* (2012) Efficient capture of *Candida albicans* and zymosan by SIGNR1 augments TLR2-dependent TNF-α production. Int Immunol. 24 (2): 89-96.
- 18. Fransen F *et al.* (2015) BALB/c and C57BL/6 Mice Differ in Polyreactive IgA Abundance, which Impacts the Generation of Antigen-Specific IgA and Microbiota Diversity. Immunity. 43 (3): 527-40.
- 19. Urso, K. *et al.* (2016) Anion Exchanger 2 Regulates Dectin-1-Dependent Phagocytosis and Killing of *Candida albicans*. PLoS One. 11 (7): e0158893.
- 20. Zhou, J. *et al.* (2016) Therapeutic targeting of myeloid-derived suppressor cells involves a novel mechanism mediated by clusterin. Sci Rep. 6: 29521.
- 21. Pinke, K.H. *et al.* (2016) Mast cells phagocyte *Candida albicans* and produce nitric oxide by mechanisms involving TLR2 and Dectin-1. <u>Immunobiology. 221 (2): 220-7.</u>

- 22. Berven, L. *et al.* (2015) Particulate yeast β-glucan is internalized by RAW 264.7 macrophages and reduces the activity of the tumor-associated protease legumain Bioactive Carbohydrates and Dietary Fibre. 6 (1): 15-23.
- 23. Walachowski, S. *et al.* (2016) Triggering Dectin-1-Pathway Alone Is Not Sufficient to Induce Cytokine Production by Murine Macrophages. PLoS One. 11 (2): e0148464.
- 24. Ferguson, B.J. *et al.* (2015) The *Schistosoma mansoni* T2 ribonuclease omega-1 modulates inflammasome-dependent IL-1β secretion in macrophages. <u>Int J Parasitol. 45</u> (13): 809-13.
- 25. Masuda, Y. *et al.* (2015) Soluble β-glucan from *Grifola frondosa* induces tumor regression in synergy with TLR9 agonist via dendritic cell-mediated immunity. <u>J Leukoc Biol. 98 (6): 1015-25.</u>
- 26. Baldwin, K.T. *et al.* (2015) Neuroinflammation triggered by β-glucan/dectin-1 signaling enables CNS axon regeneration. Proc Natl Acad Sci U S A. 112 (8): 2581-6.
- 27. Quayle K *et al.* (2015) The TLR2 agonist in polysaccharide-K is a structurally distinct lipid which acts synergistically with the protein-bound β-glucan. <u>J Nat Med. 69 (2):</u> 198-208.
- 28. Chang, T.H. *et al.* (2017) Dectin-2 is a primary receptor for NLRP3 inflammasome activation in dendritic cell response to *Histoplasma capsulatum*. <u>PLoS Pathog. 13 (7):</u> e1006485.
- 29. Seo, B.S. *et al.* (2016) Dectin-1 agonist selectively induces IgG1 class switching by LPS-activated mouse B cells. <u>Immunol Lett. 178: 114-21.</u>
- 30. Thompson, A. *et al.* (2019) The protective effect of inflammatory monocytes during systemic *C. albicans*. infection is dependent on collaboration between C-type lectin-like receptors. PLoS Pathog. 15 (6): e1007850.
- 31. Uno, A. *et al.* (2021) A novel β-glucan-oligonucleotide complex selectively delivers siRNA to APCs via Dectin-1. <u>J Control Release</u>. 338: 792-803.
- 32. Deerhake, M.E. *et al.* (2021) Dectin-1 limits autoimmune neuroinflammation and promotes myeloid cell-astrocyte crosstalk via Card9-independent expression of Oncostatin M. Immunity. 54 (3): 484-498.e8.
- 33. Niekamp, P. *et al.* (2021) Sphingomyelin Biosynthesis Is Essential for Phagocytic Signaling during *Mycobacterium tuberculosis*. Host Cell Entry. <u>mBio. 12 (1): e03141-20.</u>
- 34. Oh, S. *et al.* (2022) Pathogen size alters C-type lectin receptor signaling in dendritic cells to influence CD4 Th9 cell differentiation. <u>Cell Rep. 38 (13): 110567.</u>
- 35. Masuda, Y. *et al.* (2023) Maitake α-glucan promotes differentiation of monocytic myeloid-derived suppressor cells into M1 macrophages. <u>Life Sci. 317: 121453.</u>
- 36. Yamashita, H. *et al.* (2022) Comprehensive expression analysis with cell-type-specific transcriptome in ALS-linked mutant SOD1 mice: Revisiting the active role of glial cells in disease. Front Cell Neurosci. 16: 1045647.
- 37. Erkelens, M.N. *et al.* (2020) Intestinal Macrophages Balance Inflammatory Expression Profiles via Vitamin A and Dectin-1-Mediated Signaling. <u>Front Immunol. 11: 551.</u>
- 38. Zimara, N. *et al.* (2018) Dectin-1 Positive Dendritic Cells Expand after Infection with Leishmania major Parasites and Represent Promising Targets for Vaccine Development. Front Immunol. 9: 263.

Storage

Store at +4°C.

DO NOT FREEZE.

This product should be stored undiluted. Should this product contain a precipitate we

recommend microcentrifugation before use.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA2289 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16...)

Rabbit Anti Rat IgG (STAR17...)

<u>DyLight®800</u>

<u>FITC</u>

Goat Anti Rat IgG (STAR72...)

Goat Anti Rat IgG (STAR69...)

Goat Anti Rat IgG (STAR73...)

RPE

Rabbit Anti Rat IgG (STAR21...)

HRP

Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...) <u>DyLight®550</u>, <u>DyLight®650</u>, <u>DyLight®800</u>

Goat Anti Rat IgG (STAR131...) Alk. Phos., Biotin

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M414020:221130'

Printed on 29 Feb 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint