Datasheet: MCA2289BT

BATCH NUMBER 164961

Description:	RAT ANTI MOUSE DECTIN-1:Biotin
Specificity:	DECTIN-1
Other names:	CD369
Format:	Biotin
Product Type:	Monoclonal Antibody
Clone:	2A11
Isotype:	IgG2b
Quantity:	25 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species

Mouse

Product Form

Purified IgG conjugated to biotin - liquid

Preparation

Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

Buffer Solution

Phosphate buffered saline

Preservative Stabilisers

0.09% sodium azide (NaN_3)
1% bovine serum albumin

Approx. Protein Concentrations

IgG concentration 0.1 mg/ml

Immunogen

Dectin-1 transfected NIH3T3 cells and recombinant soluble Dectin-1.

External Database Links	**UniProt:** Q6QLQ4 Related reagents **Entrez Gene:** 56644 Clec7a Related reagents
Synonyms	Bgr, Clecsf12, Dectin1
RRID	AB_2081657
Fusion Partners	Spleen cells from immunized Fischer rats were fused with cells of the rat Y3 myeloma cell line
Specificity	**Rat anti Mouse dectin-1 antibody, clone 2A11** recognizes murine beta-glucan receptor, also known as dectin-1 or CD369. Dectin-1 is predominantly expressed by cells of the monocyte/macrophage and neutrophil lineages, but also at lower levels by dendritic cells and a subpopulation of T cells. As a major leucocyte receptor for beta-glucan this molecule may have a key role in the immunomodulatory effects of beta-glucans and in the host response to fungal pathogens. Dectin-1 may stimulate reactive oxygen production in macropahges via the protein tyrosine kinase known as Syk. Rat anti Mouse Dectin-1 antibody, clone 2A11 inhibits the binding of zymosan to macrophages via the beta-glucan receptor.
Flow Cytometry	Use 10µl of the suggested working dilution to label 10^6 cells in 100µl
References	1. Taylor, P.R. *et al.* (2002) The beta-glucan receptor, dectin-1, is predominantly expressed on the surface of cells of the monocyte/macrophage and neutrophil lineages. J Immunol. 169 (7): 3876-82. 2. Reid, D.M. *et al.* (2004) Expression of the beta-glucan receptor, Dectin-1, on murine leukocytes in situ correlates with its function in pathogen recognition and reveals potential roles in leukocyte interactions. J Leukoc Biol. 76 (1): 86-94. 3. Underhill, D.M. *et al.* (2005) Dectin-1 activates Syk tyrosine kinase in a dynamic subset of macrophages for reactive oxygen production. Blood. 106 (7): 2543-50. 4. Lefevre,L. *et al.* (2010) PPARc Ligands Switched High Fat Diet-Induced Macrophage M2b Polarization toward M2a Thereby Improving Intestinal *Candida* Elimination PLoS One. 5(9):e12828. 5. Fei, M. *et al.* (2011) TNF-alpha from inflammatory dendritic cells (DCs) regulates lung IL-17A/IL-5 levels and neutrophilia versus eosinophilia during persistent fungal infection. Proc Natl Acad Sci U S A. 108 (13): 5360-5. 6. Gazi, U. *et al.* (2011) Fungal Recognition Enhances Mannose Receptor Shedding through Dectin-1 Engagement. J Biol Chem. 286: 7822-9. 7. McDonald, J.U. *et al.* (2011) *In vivo* functional analysis and genetic modification of *in vitro*-derived mouse neutrophils. FASEB J. 25 (6): 1972-82. 8. Dewals, B.G. *et al.* (2010) IL-4Ralpha-independent expression of mannose receptor

and Ym1 by macrophages depends on their IL-10 responsiveness. PLoS Negl Trop Dis. 4: e689.
9. Galès, A. *et al.* (2010) PPARgamma controls dectin-1 expression required for host antifungal defense against *Candida albicans*. PLoS Pathog. 6: e1000714.
10. Goodridge, H.S. *et al.* (2009) Differential use of CARD9 by dectin-1 in macrophages and dendritic cells. J Immunol. 182: 1146-54.
11. Coates, P.J. *et al.* (2008) Indirect macrophage responses to ionizing radiation: implications for genotype-dependent bystander signaling. Cancer Res. 68: 450-6.
12. Dioszeghy, V. *et al.* (2008) 12/15-Lipoxygenase regulates the inflammatory response to bacterial products *in vivo*. J Immunol. 181: 6514-24.
13. Hohl, T.M. (2008) Caspofungin modulates inflammatory responses to *Aspergillus fumigatus* through stage-specific effects on fungal beta-glucan exposure. J Infect Dis. 198: 176-85.
14. Palma, A.S. *et al.* (2006) Ligands for the beta-glucan receptor, Dectin-1, assigned using "designer" microarrays of oligosaccharide probes (neoglycolipids) generated from glucan polysaccharides. J Biol Chem. 281: 5771-9.
15. Sindrilaru, A. *et al.* (2011) An unrestrained proinflammatory M1 macrophage population induced by iron impairs wound healing in humans and mice. J Clin Invest. 121: 985-97.
16. Anandasabapathy, N. *et al.* (2011) Flt3L controls the development of radiosensitive dendritic cells in the meninges and choroid plexus of the steady-state mouse brain. J Exp Med. 208 (8): 1695-705.
17. Takahara, K. *et al.* (2012) Efficient capture of *Candida albicans* and zymosan by SIGNR1 augments TLR2-dependent TNF-α production. Int Immunol. 24 (2): 89-96.
18. Fransen F *et al.* (2015) BALB/c and C57BL/6 Mice Differ in Polyreactive IgA Abundance, which Impacts the Generation of Antigen-Specific IgA and Microbiota Diversity. Immunity. 43 (3): 527-40.
19. Urso, K. *et al.* (2016) Anion Exchanger 2 Regulates Dectin-1-Dependent Phagocytosis and Killing of *Candida albicans*. PLoS One. 11 (7): e0158893.
20. Zhou, J. *et al.* (2016) Therapeutic targeting of myeloid-derived suppressor cells involves a novel mechanism mediated by clusterin. Sci Rep. 6: 29521.
21. Pinke, K.H. *et al.* (2016) Mast cells phagocyte *Candida albicans* and produce nitric oxide by mechanisms involving TLR2 and Dectin-1. Immunobiology. 221 (2): 220-7.
22. Berven, L. *et al.* (2015) Particulate yeast β-glucan is internalized by RAW 264.7 macrophages and reduces the activity of the tumor-associated protease legumain Bioactive Carbohydrates and Dietary Fibre. 6 (1): 15-23.
23. Walachowski, S. *et al.* (2016) Triggering Dectin-1-Pathway Alone Is Not Sufficient to Induce Cytokine Production by Murine Macrophages. PLoS One. 11 (2): e0148464.
24. Ferguson, B.J. *et al.* (2015) The *Schistosoma mansoni* T2 ribonuclease omega-1 modulates inflammasome-dependent IL-1β secretion in macrophages. Int J Parasitol. 45 (13): 809-13.
25. Masuda, Y. *et al.* (2015) Soluble β-glucan from *Grifola frondosa* induces tumor regression in synergy with TLR9 agonist via dendritic cell-mediated immunity. J Leukoc Biol. 98 (6): 1015-25.
26. Baldwin, K.T. *et al.* (2015) Neuroinflammation triggered by β-glucan/dectin-1 signaling enables CNS axon regeneration. Proc Natl Acad Sci U S A. 112 (8): 2581-6.
27. Quayle K *et al.* (2015) The TLR2 agonist in polysaccharide-K is a structurally distinct

lipid which acts synergistically with the protein-bound β-glucan. J Nat Med. 69 (2): 198-208.
28. Chang, T.H. *et al.* (2017) Dectin-2 is a primary receptor for NLRP3 inflammasome activation in dendritic cell response to *Histoplasma capsulatum*. PLoS Pathog. 13 (7): e1006485.
29. Seo, B.S. *et al.* (2016) Dectin-1 agonist selectively induces IgG1 class switching by LPS-activated mouse B cells. Immunol Lett. 178: 114-21.
30. Thompson, A. *et al.* (2019) The protective effect of inflammatory monocytes during systemic *C. albicans.* infection is dependent on collaboration between C-type lectin-like receptors. PLoS Pathog. 15 (6): e1007850.
31. Uno, A. *et al.* (2021) A novel β-glucan-oligonucleotide complex selectively delivers siRNA to APCs via Dectin-1. J Control Release. 338: 792-803.
32. Deerhake, M.E. *et al.* (2021) Dectin-1 limits autoimmune neuroinflammation and promotes myeloid cell-astrocyte crosstalk via Card9-independent expression of Oncostatin M. Immunity. 54 (3): 484-498.e8.
33. Niekamp, P. *et al.* (2021) Sphingomyelin Biosynthesis Is Essential for Phagocytic Signaling during *Mycobacterium tuberculosis.* Host Cell Entry. mBio. 12 (1): e03141-20.
34. Oh, S. *et al.* (2022) Pathogen size alters C-type lectin receptor signaling in dendritic cells to influence CD4 Th9 cell differentiation. Cell Rep. 38 (13): 110567.
35. Masuda, Y. *et al.* (2023) Maitake α-glucan promotes differentiation of monocytic myeloid-derived suppressor cells into M1 macrophages. Life Sci. 317: 121453.
36. Yamashita, H. *et al.* (2022) Comprehensive expression analysis with cell-type-specific transcriptome in ALS-linked mutant SOD1 mice: Revisiting the active role of glial cells in disease. Front Cell Neurosci. 16: 1045647.
37. Erkelens, M.N. *et al.* (2020) Intestinal Macrophages Balance Inflammatory Expression Profiles via Vitamin A and Dectin-1-Mediated Signaling. Front Immunol. 11: 551.
38. Zimara, N. *et al.* (2018) Dectin-1 Positive Dendritic Cells Expand after Infection with Leishmania major Parasites and Represent Promising Targets for Vaccine Development. Front Immunol. 9: 263.

Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA2289BT 10041
Regulatory	For research purposes only

Related Products

Recommended Useful Reagents

MOUSE SEROBLOCK FcR (BUF041A)
MOUSE SEROBLOCK FcR (BUF041B)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	**Worldwide**	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	**Europe**	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M414031:221130'

Printed on 27 Aug 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint