

Datasheet: MCA2290F

Description:	MOUSE ANTI HUMAN CD102:FITC			
Specificity:	CD102			
Other names:	ICAM-2			
Format:	FITC			
Product Type:	Monoclonal Antibody			
Clone:	CBRIC2/2			
Isotype:	IgG2a			
Quantity:	0.1 mg			

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	-			Neat - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human					
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid					
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)			
	FITC	490	525			
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant					
Buffer Solution	Phosphate buffered saline					
Preservative	0.09% Sodium Az	zide				
Stabilisers	1% Bovine Se	rum Albumin				
Approx. Protein Concentrations	IgG concentration	n 0.1 mg/ml				
Immunogen	Transfected COS cells expressing ICAM-2 cDNA.					
External Database Links	UniProt:					

Related reagents

P13598

Entrez Gene:

3384 ICAM2 Related reagents

RRID	AB_566413
Fusion Partners	Cells from immunised BALB/c mice were fused with cells of the Murine P3X63Ag.653 myeloma cell line.
Specificity	Mouse anti Human CD102 antibody, clone CBRIC2/2 recognizes human CD102, also known as intercellular adhesion molecule-2 (ICAM-2). CD102 is a glycosylated cell surface protein that is broadly expressed on most leucocytes and is strongly expressed by endothelial cells. CD102 interacts with the CD11a/CD18 (LFA-1). CD102 may play a role in lymphocyte re-circulation and T-cell activation (Yamashita et al. 2008).
	Mouse anti Human CD102 antibody, clone CBRIC2/2 is reported to inhibit interactions between CD102 and LFA-1 (<u>De Fougerolles <i>et al.</i> 1991</u>).
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	 Xu, H. et al. (1996) Characterization of murine intercellular adhesion molecule-2. <u>J Immunol. 156 (12): 4909-14.</u> Diacovo, T.G. et al. (1994) A functional integrin ligand on the surface of platelets: intercellular adhesion molecule-2. <u>J Clin Invest. 94 (3): 1243-51.</u> Porter, J.C. & Hall, A. (2009) Epithelial ICAM-1 and ICAM-2 regulate the egression of human T cells across the bronchial epithelium. <u>FASEB J. 23 (2): 492-502.</u> Banerjee, P. et al. (2007) Human T-cell leukemia virus type 1 (HTLV-1) p12I down-modulates ICAM-1 and -2 and reduces adherence of natural killer cells, thereby protecting HTLV-1-infected primary CD4+ T cells from autologous natural killer cell-mediated cytotoxicity despite the reduction of major histocompatibility complex class I molecules on infected cells. <u>J Virol. 81: 9707-17.</u> Arjmandi A et al. (2009) Dendritic cell adhesion to cerebral endothelium: role of endothelial cell adhesion molecules and their ligands. <u>J Neuropathol Exp Neurol. 68 (3): 300-13.</u>
Storage	Store at +4°C or at -20°C if preferred. This product should be stored undiluted. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG2a NEGATIVE CONTROL:FITC (MCA929F)

Email: antibody_sales_us@bio-rad.com

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700

Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Fax: +44 (0)1865 852 739
Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

'M366567:200529'

Printed on 11 Aug 2020

© 2020 Bio-Rad Laboratories Inc | Legal | Imprint