

# Datasheet: MCA2298PE

Description:	HAMSTER ANTI MOUSE CD29:RPE
Specificity:	CD29
Other names:	INTEGRIN BETA 1 CHAIN
Format:	RPE
<b>Product Type:</b>	Monoclonal Antibody
Clone:	HM beta 1-1
Isotype:	IgG
Quantity:	100 TESTS

### **Product Details**

## **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				Neat - 1/5

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse		
Species Cross Reactivity	reactivity is derived	from testing within our I	ons may vary between species. Cross aboratories, peer-reviewed publication ors. Please refer to references indicate
Product Form	Purified IgG conjug	ated to R. Phycoerythrir	(RPE) - lyophilized
Reconstitution	Reconstitute with 1	.0 ml distilled water	
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE 488nm laser	496	578
Preparation	Purified IgG prepare supernatant	ed by affinity chromatog	raphy on Protein G from tissue culture

Buffer Solution	Phosphate buffered saline
Preservative	0.09% sodium azide (NaN <sub>3</sub> )
Stabilisers	1% bovine serum albumin
	5% sucrose
Immunogen	Purified mouse VLA-4 antigen.
External Database	HaliPara (c
Links	UniProt:
	P09055 Related reagents
	Entrez Gene:
	16412 Itgb1 Related reagents
RRID	AB_566690
Fusion Partners	Spleen cells from immunized Armenian hamsters were fused with cells of the P3U1 mouse myeloma cell line.
Specificity	Hamster anti Mouse CD29 antibody, clone HM beta 1-1 recognizes the murine integrin beta 1 subunit (CD29), a ~110 kDa cell surface glycoprotein that is widely expressed by a variety of cells including all leucocytes. CD29 forms non-covalent bonds with the integrin alpha subunits, including CD51 and CD49a-f, to form heterodimers. The ligands for these heterodimers include collagen, fibronectin, laminin and vascular adhesion molecule-1. In the immune system beta 1 integrins play an important role in cell adhesion, migration, activation and differentiation.
	Hamster anti Mouse CD29 antibody, clone HM beta 1-1 is reported to inhibit beta 1 integrin mediated adhesion (Noto et al. 1995).
Flow Cytometry	Use 10µl of the suggested working dilution to label 10 <sup>6</sup> cells in 100µl. The Fc region of monoclonal antibodies may bind to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR (BUF041A/BUF041B).
References	<ol> <li>Eto, D.S. <i>et al.</i> (2007) Integrin-mediated host cell invasion by type 1-piliated uropathogenic Escherichia coli. <u>PLoS Pathog. 3: e100.</u></li> <li>Kouros-Mehr H (2008) GATA-3 links tumor differentiation and dissemination in a luminal</li> </ol>
	breast cancer model. <u>Cancer Cell. 13: 141-52.</u>
	3. Tiede, B.J. <i>et al.</i> (2009) A novel mouse model for non-invasive single marker tracking of mammary stem cells in vivo reveals stem cell dynamics throughout pregnancy. <u>PLoS One.</u>
	4 (11): e8035.
	4. Li S <i>et al.</i> (2010) Upregulation of CXCR4 favoring neural-like cells migration via AKT
	activation. Neurosci Res. 67 (4): 293-9.
	F. L. T.V. et al. (2010) Enithalial call adhanian malacula regulation is accordated with the

- 5. Lu, T.Y. *et al.* (2010) Epithelial cell adhesion molecule regulation is associated with the maintenance of the undifferentiated phenotype of human embryonic stem cells. <u>J Biol Chem. 285: 8719-32.</u>
- 6. Xu, L. et al. (2017) Umbilical cord-derived mesenchymal stem cells on scaffolds

facilitate collagen degradation via upregulation of MMP-9 in rat uterine scars. <u>Stem Cell</u> Res Ther. 8 (1): 84.

- 7. Hou, B. *et al.* (2018) Comparison of the Effects of BMSC-derived Schwann Cells and Autologous Schwann Cells on Remyelination Using a Rat Sciatic Nerve Defect Model. <u>Int</u> J Biol Sci. 14 (13): 1910-1922.
- 8. Porwal, K. *et al.* (2019) Increased bone marrow-specific adipogenesis by clofazimine causes impaired fracture healing, osteopenia and osteonecrosis without extra-skeletal effects in rats. Toxicol Sci. kfz172.
- 9. Zhu, C. *et al.* (2020) Antinociceptive effect of intrathecal injection of miR-9-5p modified mouse bone marrow mesenchymal stem cells on a mouse model of bone cancer pain. <u>J. Neuroinflammation</u>. 17 (1): 85.
- 10. Farahat, M. *et al.* (2022) Mechanotransductive Mechanisms of Biomimetic Hydrogel Cues Modulating Meckel's Cartilage Degeneration. Adv Biol (Weinh). 2022: e2101315.
- 11. Zbinden, A. *et al.* (2021) Nidogen-1 Mitigates Ischemia and Promotes Tissue Survival and Regeneration. <u>Adv Sci (Weinh)</u>. 8 (4): 2002500.
- 12. Chaker, D. *et al.* (2024) Direct Reprogramming of Hepatocytes Into JAK/Stat-Dependent LGR5+ Liver Cells Able to Initiate Intrahepatic Cholangiocarcinoma. <u>Stem Cells. 42 (4): 301-316.</u>

Storage	This product is shipped at ambient temperature
	Tillo product lo chippod at ambient temperatare

Store at +4°C.

DO NOT FREEZE

This product should be stored undiluted. This product is photosensitive and should be protected from light.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #20487 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA2298PE">https://www.bio-rad-antibodies.com/SDS/MCA2298PE</a> 20487

**Regulatory** For research purposes only

### Related Products

#### **Recommended Negative Controls**

HAMSTER (ARMENIAN) IgG NEGATIVE CONTROL:RPE (MCA2356PE)

#### **Recommended Positive Controls**

HAMSTER (ARMENIAN) IgG NEGATIVE CONTROL:RPE (MCA2356PE)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M440530:250523'

Printed on 23 May 2025