

Datasheet: MCA2305GA

Description:	RAT ANTI MOUSE CD16/CD32
Specificity:	CD16/CD32
Other names:	MOUSE SEROBLOCK FcR
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	FCR4G8
Isotype:	IgG2b
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/25 - 1/50
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% sodium azide (NaN ₃)

Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	PU5 1.8 IOE7 Balb/c mouse cell line.
External Database Links	<p>UniProt:</p> <p>P08508 Related reagents</p> <p>P08101 Related reagents</p> <p>Entrez Gene:</p> <p>14131 Fcgr3 Related reagents</p> <p>14130 Fcgr2b Related reagents</p>
Synonyms	Fcgr2b, Ly-17
RRID	AB_2262717
Fusion Partners	Cells from immunised rats were fused with cells of the mouse P3X63-Ag8.653 myeloma cell line.
Specificity	Rat anti Mouse CD16/CD32 antibody, clone FCR4G8 recognizes an epitope expressed by the murine low affinity Fc receptors, Fc gamma III (CD16) and Fc gamma II (CD32). In the mouse only the transmembrane form of CD16 is reported to exist, which is expressed on macrophages, NK cells, granulocytes, myeloid precursors, and subpopulations of T lymphocytes. CD32 is primarily expressed on cells of the myeloid lineage and also on mature B lymphocytes.
Flow Cytometry	Use 10µl of the suggested working dilution to label 10 ⁶ cells in 100µl
References	<ol style="list-style-type: none"> Hart, R. & Greaves, D.R. (2010) Chemerin contributes to inflammation by promoting macrophage adhesion to VCAM-1 and fibronectin through clustering of VLA-4 and VLA-5. J Immunol. 185 (6): 3728-39. da Silva, R.G. <i>et al.</i> (2010) Endothelial alpha3beta1-integrin represses pathological angiogenesis and sustains endothelial-VEGF. Am J Pathol. 177 (3): 1534-48. Birjandi, S.Z. <i>et al.</i> (2011) Alterations in marginal zone macrophages and marginal zone B cells in old mice. J Immunol. 186 (6): 3441-51. Lunnon, K. <i>et al.</i> (2011) Systemic inflammation modulates Fc receptor expression on microglia during chronic neurodegeneration. J Immunol. 186 (12): 7215-24. Birjandi, S.Z. <i>et al.</i> (2011) Alterations in marginal zone macrophages and marginal zone B cells in old mice. J Immunol. 186: 3441-51. Hart, A.D. <i>et al.</i> (2012) Age related changes in microglial phenotype vary between CNS regions: grey versus white matter differences. Brain Behav Immun. 26 (5): 754-65. Teeling, J.L. <i>et al.</i> (2012) Intracerebral immune complex formation induces inflammation in the brain that depends on Fc receptor interaction. Acta Neuropathol. 124 (4): 479-90. Jones, D.T. <i>et al.</i> (2013) Endogenous ribosomal protein L29 (RPL29): a newly identified

regulator of angiogenesis in mice. [Dis Model Mech. 6: 115-24.](#)

9. Murinello, S. *et al.* (2014) Fcγ receptor upregulation is associated with immune complex inflammation in the mouse retina and early age-related macular degeneration. [Invest Ophthalmol Vis Sci. 55 \(1\): 247-58.](#)

10. Kapellos, T.S. *et al.* (2016) A novel real time imaging platform to quantify macrophage phagocytosis. [Biochem Pharmacol. 116: 107-19.](#)

11. Ahmed, N. *et al.* (2022) Blood stem cell PU.1 upregulation is a consequence of differentiation without fast autoregulation. [J Exp Med. 219 \(1\): e20202490.](#)

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA2305GA>
10040

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Rat IgG (STAR69...)

[FITC](#)

Goat Anti Rat IgG (STAR73...)

[RPE](#)

Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)

[DyLight@550](#), [DyLight@650](#), [DyLight@800](#)

Rabbit Anti Rat IgG (STAR21...)

[HRP](#)

Rabbit Anti Rat IgG (STAR16...)

[DyLight@800](#)

Goat Anti Rat IgG (STAR131...)

[Alk. Phos.](#), [Biotin](#)

Rabbit Anti Rat IgG (STAR17...)

[FITC](#)

Goat Anti Rat IgG (STAR72...)

[HRP](#)

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751
Email: antibody_sales_us@bio-rad.com

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739
Email: antibody_sales_uk@bio-rad.com

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50
Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M414326:221206'

Printed on 18 Jan 2024