

Datasheet: MCA2307F

Description:	MOUSE ANTI PIG CD5:FITC
Specificity:	CD5
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	1H6/8
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10
Immunoprecipitation	▪			
Western Blotting	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Pig		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide		
Stabilisers	1% Bovine Serum Albumin		
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml		

Immunogen	ConA/PMA activated porcine peripheral blood cells.
External Database Links	UniProt: Q9GMA0 Related reagents
RRID	AB_566839
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse X63-Ag.8.653 myeloma cell line.
Specificity	<p>Mouse anti Pig CD5 antibody, clone 1H6/8 recognizes pig CD5. Clone 1H6/8 was clustered as CD5a at the Second International Swine CD antigen Workshop (Saalmuller et al. 1998), based on its cellular distribution, ability to completely inhibit the binding of monoclonal antibody clone b53b7, the internal control for the second workshop and the immunoprecipitation of the same 55 kDa + 60 kDa doublet as the control antibody (Pescowitz et al. 1998). Porcine CD5 is expressed by all circulating peripheral blood T lymphocytes and some B cells. CD5 is not expressed by porcine γ/δ T cells (Saalmuller et al. 1994) and can hence be used for the discrimination between NK cells (CD4- CD8+ CD5-) and MHC-restricted cytotoxic T cells (CD4- CD8+ CD5+).</p> <p>No reactivity of Mouse anti Pig CD5 antibody, clone 1H6/8 was noted with canine, bovine, equine or human tissue samples (Doménech et al. 2003).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to 1×10^6 cells in 100ul.
References	<ol style="list-style-type: none"> Pescovitz, M.D. <i>et al.</i> (1998) Analyses of monoclonal antibodies reacting with porcine CD5: results from the Second International Swine CD Workshop. Vet Immunol Immunopathol. 60 (3-4): 269-73. Saalmüller, A. <i>et al.</i> (1998) Overview of the Second International Workshop to define swine cluster of differentiation (CD) antigens. Vet Immunol Immunopathol. 60 (3-4): 207-28. Doménech, N. <i>et al.</i> (2003) A new epitope on swine CD5 molecule detected by monoclonal antibody 5F12/9. Hybrid Hybridomics. 22 (3): 179-82. Sánchez-Torres, C. <i>et al.</i> (2003) Expression of porcine CD163 on monocytes/macrophages correlates with permissiveness to African swine fever infection. Arch Virol. 148 (12): 2307-23. Forner, R. <i>et al.</i> (2021) Distribution difference of colostrum-derived B and T cells subsets in gilts and sows. PLoS One. 16 (5): e0249366. Skovdal, S.M. <i>et al.</i> (2019) Inhaled nebulized glatiramer acetate against Gram-negative bacteria is not associated with adverse pulmonary reactions in healthy, young adult female pigs. PLoS One. 14 (10): e0223647. Jarosz, Ł. <i>et al.</i> (2021) The Effect of Feed Supplementation with EM Bokashi® Multimicrobial Probiotic Preparation on Selected Parameters of Sow Colostrum and Milk as Indicators of the Specific and Nonspecific Immune Response. Probiotics Antimicrob Proteins. Oct 01 [Epub ahead of print].
Further Reading	1. Piriou-Guzylack, L. (2008) Membrane markers of the immune cells in swine: an update.

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10041 available at: 10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL:FITC \(MCA929F\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M385152:210513'

Printed on 21 Mar 2022