

Datasheet: MCA2312F

BATCH NUMBER 174203

Description:	MOUSE ANTI PIG CD172a:FITC
Specificity:	CD172a
Other names:	SWC3
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	BL1H7
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Pig		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% sodium azide (NaN ₃)		
Stabilisers	1% bovine serum albumin		
Approx. Protein	IgG concentration 0.1 mg/ml		

Concentrations

Immunogen Porcine alveolar macrophages.

External Database Links

UniProt:

[Q5K4Q3](#)

[Related reagents](#)

RRID AB_2188073

Fusion Partners Spleen cells from immunised BALB/c mice were fused with cells of the mouse SP2/0 myeloma cell line.

Specificity **Mouse anti Pig CD172a, clone BL1H7** recognizes porcine CD172a, a member of the signal regulatory protein (SIRP) family ([Alvarez et al. 2000](#)).

Mouse anti Pig CD172a, clone BL1H7 was originally clustered as SWC3 at the Third International Swine Cluster of Differentiation Workshop ([Haverson et al. 2001](#); [Thacker et al. 2001](#)). CD172a is expressed on monocyte derived dendritic cells (MoDCs) ([Facci et al. 2010](#)) also conventional (cDCs), plasmacytoid (pDCs) DCs and blood DCs. ([Facci](#); [Jeong et al. 2010](#)). Mouse anti Pig CD172a, clone BL1H7 immunoprecipitates a single band of ~90-110 kDa from preparations of biotinylated alveolar macrophages, a result confirmed by Western blotting analysis of alveolar macrophage lysates under non reducing conditions ([Alvarez et al. 2000](#)). Aberrant expression of CD172a has been noted on porcine leukemias ([Sipos et al. 2006](#)) with blast cells co-expressing lymphocytic markers CD5 and CD25 whilst expressing the Myeloid marker CD172a in a [bi-phenotypic pattern](#) as opposed to the more characteristic [single population](#) of CD172+ cells seen in normal blood PBMC ([Chamorro et al. 2005](#)).

Mouse anti Pig CD172a, clone BL1H7 has proved a useful and reliable tool for immunohistochemical analysis of routinely processed, formalin fixed, paraffin embedded porcine tissues ([Domenech et al. 2003](#)).

Flow Cytometry Use 10µl of the suggested working dilution to 1x10⁶ cells in 100µl

References

1. Alvarez, B. *et al.* (2000) A porcine cell surface receptor identified by monoclonal antibodies to SWC3 is a member of the signal regulatory protein family and associates with protein-tyrosine phosphatase SHP-1. [Tissue Antigens. 55 \(4\): 342-51.](#)
2. Domenech, N. *et al.* (2003) Identification of porcine macrophages with monoclonal antibodies in formalin-fixed, paraffin-embedded tissues. [Vet Immunol Immunopathol. 94 \(1-2\): 77-81.](#)
3. Carrillo, A. *et al.* (2002) Isolation and characterization of immortalized porcine aortic endothelial cell lines. [Vet Immunol Immunopathol. 89 \(1-2\): 91-8.](#)
4. Fraile, L. *et al.* (2012) Immunomodulatory properties of beta-sitosterol in pig immune responses. [Int Immunopharmacol. 13 \(3\): 316-21.](#)
5. Jeong, H.J. *et al.* (2010) Comparative measurement of cell-mediated immune responses of swine to the M and N proteins of porcine reproductive and respiratory syndrome virus. [Clin Vaccine Immunol. 17 \(4\): 503-12.](#)

6. Gimeno, M. *et al.* (2011) Cytokine profiles and phenotype regulation of antigen presenting cells by genotype-I porcine reproductive and respiratory syndrome virus isolates. [Vet Res. 42: 9.](#)
7. Moreno, S. *et al.* (2010) Porcine monocyte subsets differ in the expression of CCR2 and in their responsiveness to CCL2. [Vet Res. 41: 76.](#)
8. Facci, M.R. *et al.* (2010) A comparison between isolated blood dendritic cells and monocyte-derived dendritic cells in pigs. [Immunology. 129: 396-405.](#)
9. Clapperton, M. *et al.* (2005) Innate immune traits differ between Meishan and Large White pigs. [Vet Immunol Immunopathol. 104: 131-44.](#)
10. Argilaguet, J.M. *et al.* (2012) DNA vaccination partially protects against African swine fever virus lethal challenge in the absence of antibodies. [PLoS One. 7 \(9\): e40942.](#)
11. Kapetanovic, R. *et al.* (2012) Pig bone marrow-derived macrophages resemble human macrophages in their response to bacterial lipopolysaccharide. [J Immunol. 188: 3382-94.](#)
12. Tambuyzer, B.R. *et al.* (2012) Osteopontin alters the functional profile of porcine microglia *in vitro*. [Cell Biol Int. 36 \(12\): 1233-8.](#)
13. Robinson, S.R. *et al.* (2015) Broadly neutralizing antibodies against the rapidly evolving porcine reproductive and respiratory syndrome virus. [Virus Res. 203: 56-65.](#)
14. Li, J. & Murtaugh, M.P. (2015) Functional analysis of porcine reproductive and respiratory syndrome virus N-glycans in infection of permissive cells. [Virology. 477: 82-8.](#)
15. Prims, S. *et al.* (2016) Intestinal immune cell quantification and gram type classification of the adherent microbiota in conventionally and artificially reared, normal and low birth weight piglets. [J Livestock Sci 185: 1-7.](#)
16. Gardner, D.S. *et al.* (2016) Remote effects of acute kidney injury in a porcine model. [Am J Physiol Renal Physiol. 310 \(4\): F259-71.](#)
17. Valekova I *et al.* (2016) Revelation of the IFN α , IL-10, IL-8 and IL-1 β as promising biomarkers reflecting immuno-pathological mechanisms in porcine Huntington's disease model. [J Neuroimmunol. 293: 71-81.](#)
18. Gardner, D.S. *et al.* (2016) Remote effects of acute kidney injury in a porcine model. [Am J Physiol Renal Physiol. 310 \(4\): F259-71.](#)
19. Thirion-Delalande, C. *et al.* (2017) Comparative analysis of the oral mucosae from rodents and non-rodents: Application to the nonclinical evaluation of sublingual immunotherapy products. [PLoS One. 12 \(9\): e0183398.](#)
20. Auray, G. *et al.* (2013) Porcine neonatal blood dendritic cells, but not monocytes, are more responsive to TLRs stimulation than their adult counterparts. [PLoS One. 8 \(5\): e59629.](#)
21. Hu, W. *et al.* (2021) shRNA transgenic swine display resistance to infection with the foot-and-mouth disease virus. [Sci Rep. 11 \(1\): 16377.](#)
22. Radlowski, E.C. *et al.* (2021) Combination-Feeding Causes Differences in Aspects of Systemic and Mucosal Immune Cell Phenotypes and Functions Compared to Exclusive Sow-Rearing or Formula-Feeding in Piglets. [Nutrients. 13\(4\):1097.](#)
23. Melgoza-González, A.E. *et al.* (2022) Antigen Targeting of Porcine Skin DEC205+ Dendritic Cells [Vaccines. 10 \(5\): 684.](#)
24. Zhou, L. *et al.* (2022) Clinical improvement of sepsis by extracorporeal centrifugal leukocyte apheresis in a porcine model. [J Transl Med. 20 \(1\): 538.](#)
25. Álvarez, B. *et al.* (2023) Porcine Macrophage Markers and Populations: An Update. [Cells. 12 \(16\): 2103.](#)
26. Wei, H. *et al.* (2024) Novel Approach in DNA vaccine development against porcine

circovirus type 2 [Theoretical and Applied Veterinary Medicine. 12 \(1\): 15-24.](#)

27. Chen, Y. *et al.* (2025) ASFV activates STAT3 to induce proviral M2 macrophage polarization. [Vet Microbiol. 311: 110733.](#)

28. Jaudas, F. *et al.* (2025) Perinatal dysfunction of innate immunity in cystic fibrosis. [Sci Transl Med. 17 \(782\): eadk9145.](#)

Further Reading	1. Piriou-Guzylack, L. (2008) Membrane markers of the immune cells in swine: an update. Vet Res. 39: 54.
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA2312F
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M414391:221206'

Printed on 21 Aug 2026