

Datasheet: MCA2312GA

BATCH NUMBER 148204

Description:	MOUSE ANTI PIG CD172a
Specificity:	CD172a
Other names:	SWC3
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	BL1H7
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10 - 1/20
Immunohistology - Frozen	▪			
Immunohistology - Paraffin (1)	▪			
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting (2)	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1) This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose.

(2) Clone BL1H7 recognizes porcine CD172a under non-reducing conditions.

Target Species	Pig
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline

Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Porcine alveolar macrophages.
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse SP2/0 myeloma cell line.
Specificity	<p>Mouse anti Pig CD172a, clone BL1H7 recognizes porcine CD172a, a member of the signal regulatory protein (SIRP) family (Alvarez et al. 2000). Mouse anti Pig CD172a, clone BL1H7 was originally clustered as SWC3 at the Third International Swine Cluster of Differentiation Workshop (Haverson et al. 2001; Thacker et al. 2001). CD172a is expressed on monocyte derived dendritic cells (MoDCs) (Facci et al. 2010) also conventional (cDCs), plasmacytoid (pDCs) DCs and blood DCs. (Facci; Jeong et al. 2010). Mouse anti Pig CD172a, clone BL1H7 immunoprecipitates a single band of ~90-110 kDa from preparations of biotinylated alveolar macrophages, a result confirmed by Western blotting analysis of alveolar macrophage lysates under non reducing conditions (Alvarez et al. 2000).</p> <p>Mouse anti Pig CD172a, clone BL1H7 has proved a useful and reliable tool for immunohistochemical analysis of routinely processed, formalin fixed, paraffin embedded porcine tissues (Domenech et al. 2003).</p> <p>Aberrant expression of CD172a has been noted on porcine leukemias (Sipos et al. 2006) with blast cells co-expressing lymphocytic markers CD5 and CD25 whilst expressing the Myeloid marker CD172a in a bi-phenotypic pattern as opposed to the more characteristic single population of CD172+ cells seen in normal blood PBMC (Chamorro et al. 2005).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to 1x10 ⁶ cells in 100ul.
Histology Positive Control Tissue	Porcine spleen.
Western Blotting	MCA2312GA detects a band of approximately 90-115 kDa in alveolar macrophage lysates.
References	<ol style="list-style-type: none"> 1. Alvarez, B. <i>et al.</i> (2000) A porcine cell surface receptor identified by monoclonal antibodies to SWC3 is a member of the signal regulatory protein family and associates with protein-tyrosine phosphatase SHP-1. Tissue Antigens. 55 (4): 342-51. 2. Domenech, N. <i>et al.</i> (2003) Identification of porcine macrophages with monoclonal antibodies in formalin-fixed, paraffin-embedded tissues. Vet Immunol Immunopathol. 94 (1-2): 77-81. 3. Carrillo, A. <i>et al.</i> (2002) Isolation and characterization of immortalized porcine aortic endothelial cell lines. Vet Immunol Immunopathol. 89 (1-2): 91-8.

4. Fraile, L. *et al.* (2012) Immunomodulatory properties of beta-sitosterol in pig immune responses. [Int Immunopharmacol. 13 \(3\): 316-21.](#)
5. Jeong, H.J. *et al.* (2010) Comparative measurement of cell-mediated immune responses of swine to the M and N proteins of porcine reproductive and respiratory syndrome virus. [Clin Vaccine Immunol. 17 \(4\): 503-12.](#)
6. Gimeno, M. *et al.* (2011) Cytokine profiles and phenotype regulation of antigen presenting cells by genotype-I porcine reproductive and respiratory syndrome virus isolates. [Vet Res. 42: 9.](#)
7. Moreno, S. *et al.* (2010) Porcine monocyte subsets differ in the expression of CCR2 and in their responsiveness to CCL2. [Vet Res. 41: 76.](#)
8. Facci, M.R. *et al.* (2010) A comparison between isolated blood dendritic cells and monocyte-derived dendritic cells in pigs. [Immunology. 129: 396-405.](#)
9. Clapperton, M. *et al.* (2005) Innate immune traits differ between Meishan and Large White pigs. [Vet Immunol Immunopathol. 104: 131-44.](#)
10. Argilaguet, J.M. *et al.* (2012) DNA vaccination partially protects against African swine fever virus lethal challenge in the absence of antibodies. [PLoS One. 7 \(9\): e40942.](#)
11. Kapetanovic, R. *et al.* (2012) Pig bone marrow-derived macrophages resemble human macrophages in their response to bacterial lipopolysaccharide. [J Immunol. 188: 3382-94.](#)
12. Tambuyzer, B.R. *et al.* (2012) Osteopontin alters the functional profile of porcine microglia *in vitro*. [Cell Biol Int. 36 \(12\): 1233-8.](#)
13. Robinson, S.R. *et al.* (2015) Broadly neutralizing antibodies against the rapidly evolving porcine reproductive and respiratory syndrome virus. [Virus Res. 203: 56-65.](#)
14. Li, J. & Murtaugh, M.P. (2015) Functional analysis of porcine reproductive and respiratory syndrome virus N-glycans in infection of permissive cells. [Virology. 477: 82-8.](#)
15. Prims, S. *et al.* (2016) Intestinal immune cell quantification and gram type classification of the adherent microbiota in conventionally and artificially reared, normal and low birth weight piglets. [J Livestock Sci 185: 1-7.](#)
16. Gardner, D.S. *et al.* (2016) Remote effects of acute kidney injury in a porcine model. [Am J Physiol Renal Physiol. 310 \(4\): F259-71.](#)
17. Valekova I *et al.* (2016) Revelation of the IFN α , IL-10, IL-8 and IL-1 β as promising biomarkers reflecting immuno-pathological mechanisms in porcine Huntington's disease model. [J Neuroimmunol. 293: 71-81.](#)
18. Gardner, D.S. *et al.* (2016) Remote effects of acute kidney injury in a porcine model. [Am J Physiol Renal Physiol. 310 \(4\): F259-71.](#)
19. Thirion-Delalande, C. *et al.* (2017) Comparative analysis of the oral mucosae from rodents and non-rodents: Application to the nonclinical evaluation of sublingual immunotherapy products. [PLoS One. 12 \(9\): e0183398.](#)
20. Auray, G. *et al.* (2013) Porcine neonatal blood dendritic cells, but not monocytes, are more responsive to TLRs stimulation than their adult counterparts. [PLoS One. 8 \(5\): e59629.](#)

Further Reading

1. Piriou-Guzylack, L. (2008) Membrane markers of the immune cells in swine: an update. [Vet Res. 39: 54.](#)

Storage

Store at +4°C or at -20°C if preferred.
 Storage in frost-free freezers is not recommended.
 This product should be stored undiluted. Avoid repeated freezing and thawing as this may

denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA2312GA 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (STAR70...)	FITC
Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M366627:200529'

Printed on 12 Mar 2025