

Datasheet: MCA2314F

BATCH NUMBER 171190

Description:	MOUSE ANTI PIG SLA CLASS II DR:FITC
Specificity:	SLA CLASS II DR
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	2E9/13
Isotype:	IgG2b
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species

Pig

Species Cross Reactivity

Reacts with: Bovine

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid

Max Ex/Em

Fluorophore	Excitation Max (nm)	Emission Max (nm)
FITC	490	525

Preparation

Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant

Buffer Solution

Phosphate buffered saline

Preservative	0.09% sodium azide (NaN ₃)
Stabilisers	1% bovine serum albumin
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml
Immunogen	Porcine monocytes.
External Database Links	UniProt: Q85ZW4 Related reagents
RRID	AB_567376
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the mouse X63-Ag.8.653 myeloma cell line.
Specificity	<p>Mouse anti Pig SLA Class II DR antibody, clone 2E9/13 recognizes SLA DR molecules which are expressed on all B cells, antigen presenting cells and on certain subsets of resting and activated T cells. Mouse anti Pig SLA Class II DR antibody, clone 289/13 reacts with lymphocytes from all outbred and miniature pigs so far tested, suggesting that it recognizes a monomorphic determinant of porcine SLA DR.</p> <p>The major histocompatibility complex (MHC) is a cluster of genes that are important in the immune response to infections. In pigs, this is referred to as the swine leukocyte antigen (SLA) region. There are 3 major MHC class II proteins encoded by the SLA which are SLA DP, SLA DQ and SLA DR.</p> <p>Mouse anti pig SLA class II DR, clone 2E9/13 immunoprecipitates a heterodimer composed of two polypeptides of ~28 and ~35 kDa from NP-40 extracts of biotin surface-labeled porcine peripheral blood mononuclear cells. Mouse anti Pig SLA Class II DR antibody, clone 289/13 is reported to inhibit the mixed lymphocyte reaction and T cell stimulation induced by African swine fever virus and staphylococcal enterotoxin B (Bullido <i>et al.</i> 1997).</p>
Flow Cytometry	Use 10µl of the suggested working dilution to 1x10 ⁶ cells in 100µl
References	<ol style="list-style-type: none"> 1. Bullido, R. <i>et al.</i> (1997) Characterization of five monoclonal antibodies specific for swine class II major histocompatibility antigens and crossreactivity studies with leukocytes of domestic animals. Dev Comp Immunol. 21 (3): 311-22. 2. Jeong, H.J. <i>et al.</i> (2010) Comparative measurement of cell-mediated immune responses of swine to the M and N proteins of porcine reproductive and respiratory syndrome virus. Clin Vaccine Immunol. 17: 503-12. 3. Ding, Q. <i>et al.</i> (2011) Human PD-L1-overexpressing porcine vascular endothelial cells induce functionally suppressive human CD4+CD25hiFoxp3+ Treg cells. J Leukoc Biol. 90 (1): 77-86. 4. Thierry, A. <i>et al.</i> (2012) Identification of invariant natural killer T cells in porcine peripheral blood. Vet Immunol Immunopathol. 149 (3-4): 272-9.

5. Iwase H *et al.* (2015) Initial *in vivo* experience of pig artery patch transplantation in baboons using mutant MHC (CIITA-DN) pigs. [Transpl Immunol. 32 \(2\): 99-108.](#)
6. Zanotti, C. *et al.* (2015) Differential Biological Activities of Swine Interferon- α Subtypes. [J Interferon Cytokine Res. 35 \(12\): 990-1002.](#)
7. Wang, Y. *et al.* (2016) Genipin crosslinking reduced the immunogenicity of xenogeneic decellularized porcine whole-liver matrices through regulation of immune cell proliferation and polarization. [Sci Rep. 6: 24779.](#)
8. Gardner, D.S. *et al.* (2016) Remote effects of acute kidney injury in a porcine model. [Am J Physiol Renal Physiol. 310 \(4\): F259-71.](#)
9. Singleton, H. *et al.* (2016) Establishing Porcine Monocyte-Derived Macrophage and Dendritic Cell Systems for Studying the Interaction with PRRSV-1. [Front Microbiol. 7: 832.](#)
10. Rahe, M.C. & Murtaugh, M.P. (2017) Interleukin-21 Drives Proliferation and Differentiation of Porcine Memory B Cells into Antibody Secreting Cells. [PLoS One. 12 \(1\): e0171171.](#)
11. Mašek, J. *et al.* (2017) Multi-layered nanofibrous mucoadhesive films for buccal and sublingual administration of drug-delivery and vaccination nanoparticles - important step towards effective mucosal vaccines. [J Control Release. 249: 183-95.](#)
12. Ladowski, J.M. *et al.* (2018) Swine Leukocyte Antigen Class II Is a Xenoantigen. [Transplantation. 102 \(2\): 249-54.](#)
13. Yang, N. *et al.* (2018) Reduced antigen presentation capability and modified inflammatory/immunosuppressive cytokine expression of induced monocyte-derived dendritic cells from peripheral blood of piglets infected with porcine circovirus type 2. [Arch Virol. 163 \(5\): 1231-9.](#)
14. López, E. *et al.* (2019) Identification of very early inflammatory markers in a porcine myocardial infarction model. [BMC Vet Res. 15 \(1\): 91.](#)
15. Liu, S. *et al.* (2019) Endothelial IL-8 induced by porcine circovirus type 2 affects dendritic cell maturation and antigen-presenting function. [Virol J. 16 \(1\): 154.](#)
16. Radlowski, E.C. *et al.* (2021) Combination-Feeding Causes Differences in Aspects of Systemic and Mucosal Immune Cell Phenotypes and Functions Compared to Exclusive Sow-Rearing or Formula-Feeding in Piglets. [Nutrients. 13\(4\):1097.](#)
17. Arenal, Á. *et al.* (2022) Effects of Cardiac Stem Cell on Postinfarction Arrhythmogenic Substrate. [Int J Mol Sci. 23 \(24\): 16211.](#)
18. Franzoni, G. *et al.* (2022) Analyses of the Impact of Immunosuppressive Cytokines on Porcine Macrophage Responses and Susceptibility to Infection to African Swine Fever Viruses. [Pathogens. 11 \(2\): 166.](#)
19. Haach, V. *et al.* (2023) A polyvalent virosomal influenza vaccine induces broad cellular and humoral immunity in pigs. [Virol J. 20 \(1\): 181.](#)
20. Skirecki, T. *et al.* (2022) Compartment-Specific Differences in the Activation of Monocyte Subpopulations Are Not Affected by Nitric Oxide and Glucocorticoid Treatment in a Model of Resuscitated Porcine Endotoxemic Shock. [J Clin Med. 11 \(9\): 2641.](#)
21. Yuan, C. *et al.* (2024) Comparison of B cells' immune response induced by PEDV virulent and attenuated strains. [Front Microbiol. 15: 1344344.](#)
22. Zhang, M. *et al.* (2024) PCV2 Induced Endothelial Derived IL-8 Affects MoDCs Maturation Mainly via NF- κ B Signaling Pathway. [Viruses. 16 \(4\): 646.](#)
23. Tng, P.Y.L. *et al.* (2025) Early disruption of the innate-adaptive immune axis *in vivo* after infection with virulent Georgia 2007/1 ASFV. [Discov Immunol. 4 \(1\): kyaf014.](#)

- Further Reading**
1. Piriou-Guzylack, L. (2008) Membrane markers of the immune cells in swine: an update. [Vet Res. 39: 54.](#)
 2. Rayat GR *et al.* (2016) First update of the International Xenotransplantation Association consensus statement on conditions for undertaking clinical trials of porcine islet products in type 1 diabetes - Chapter 3: Porcine islet product manufacturing and release testing criteria. [Xenotransplantation. 23 \(1\): 38-45.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10041 available at: <https://www.bio-rad-antibodies.com/SDS/MCA2314F>

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG2b NEGATIVE CONTROL:FITC \(MCA691F\)](#)

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M437789:250319'

Printed on 06 Feb 2026