

# Datasheet: MCA2314GA

**BATCH NUMBER 161441**

<b>Description:</b>	MOUSE ANTI PIG SLA CLASS II DR
<b>Specificity:</b>	SLA CLASS II DR
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	2E9/13
<b>Isotype:</b>	IgG2b
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/25 - 1/200
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Pig
<b>Species Cross Reactivity</b>	<p>Reacts with: Bovine</p> <p><b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.</p>
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN <sub>3</sub> )
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Porcine monocytes.
External Database Links	<b>UniProt:</b> <a href="#">Q85ZW4</a> <a href="#">Related reagents</a>
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse X63-Ag.8.653 myeloma cell line.
Specificity	<p><b>Mouse anti Pig SLA Class II DR antibody, clone 2E9/13</b> recognizes SLA DR molecules which are expressed on all B cells, antigen presenting cells and on certain subsets of resting and activated T cells. Mouse anti Pig SLA Class II DR antibody, clone 289/13 reacts with lymphocytes from all outbred and miniature pigs so far tested, suggesting that it recognizes a monomorphic determinant of porcine SLA DR.</p> <p>The major histocompatibility complex (MHC) is a cluster of genes that are important in the immune response to infections. In pigs, this is referred to as the swine leukocyte antigen (SLA) region. There are 3 major MHC class II proteins encoded by the SLA which are SLA DP, SLA DQ and SLA DR.</p> <p>Mouse anti pig SLA class II DR, clone 2E9/13 immunoprecipitates a heterodimer composed of two polypeptides of ~28 and ~35 kDa from NP-40 extracts of biotin surface-labeled porcine peripheral blood mononuclear cells. Mouse anti Pig SLA Class II DR antibody, clone 289/13 is reported to inhibit the mixed lymphocyte reaction and T cell stimulation induced by African swine fever virus and staphylococcal enterotoxin B (<a href="#">Bullido et al. 1997</a>).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to 1x10 <sup>6</sup> cells in 100ul.
References	<ol style="list-style-type: none"> <li>1. Bullido, R. et al. (1997) Characterization of five monoclonal antibodies specific for swine class II major histocompatibility antigens and crossreactivity studies with leukocytes of domestic animals. <a href="#">Dev Comp Immunol. 21 (3): 311-22.</a></li> <li>2. Jeong, H.J. et al. (2010) Comparative measurement of cell-mediated immune responses of swine to the M and N proteins of porcine reproductive and respiratory syndrome virus. <a href="#">Clin Vaccine Immunol. 17: 503-12.</a></li> <li>3. Ding, Q. et al. (2011) Human PD-L1-overexpressing porcine vascular endothelial cells induce functionally suppressive human CD4+CD25hiFoxp3+ Treg cells. <a href="#">J Leukoc Biol. 90 (1): 77-86.</a></li> <li>4. Wang, Y. et al. (2016) Genipin crosslinking reduced the immunogenicity of xenogeneic</li> </ol>

decellularized porcine whole-liver matrices through regulation of immune cell proliferation and polarization. [Sci Rep. 6: 24779.](#)

5. Park KM *et al.* (2013) Generation of porcine induced pluripotent stem cells and evaluation of their major histocompatibility complex protein expression in vitro. [Vet Res Commun. 37 \(4\): 293-301.](#)

6. Iwase H *et al.* (2015) Initial *in vivo* experience of pig artery patch transplantation in baboons using mutant MHC (CIITA-DN) pigs. [Transpl Immunol. 32 \(2\): 99-108.](#)

7. Singleton, H. *et al.* (2016) Establishing Porcine Monocyte-Derived Macrophage and Dendritic Cell Systems for Studying the Interaction with PRRSV-1. [Front Microbiol. 7: 832.](#)

8. Zanutti, C. *et al.* (2015) Differential Biological Activities of Swine Interferon- $\alpha$  Subtypes. [J Interferon Cytokine Res. 35 \(12\): 990-1002.](#)

9. Rayat GR *et al.* (2016) First update of the International Xenotransplantation Association consensus statement on conditions for undertaking clinical trials of porcine islet products in type 1 diabetes - Chapter 3: Porcine islet product manufacturing and release testing criteria. [Xenotransplantation. 23 \(1\): 38-45.](#)

10. Mašek J *et al.* (2016) Multi-layered nanofibrous mucoadhesive films for buccal and sublingual administration of drug-delivery and vaccination nanoparticles - important step towards effective mucosal vaccines. [J Control Release. Jul 25. pii: S0168-3659\(16\)30471-0 \[Epub ahead of print\]](#)

11. Gardner, D.S. *et al.* (2016) Remote effects of acute kidney injury in a porcine model. [Am J Physiol Renal Physiol. 310 \(4\): F259-71.](#)

12. Rahe, M.C. & Murtaugh, M.P. (2017) Interleukin-21 Drives Proliferation and Differentiation of Porcine Memory B Cells into Antibody Secreting Cells. [PLoS One. 12 \(1\): e0171171.](#)

13. López, E. *et al.* (2019) Identification of very early inflammatory markers in a porcine myocardial infarction model. [BMC Vet Res. 15 \(1\): 91.](#)

14. Yang, N. *et al.* (2018) Reduced antigen presentation capability and modified inflammatory/immunosuppressive cytokine expression of induced monocyte-derived dendritic cells from peripheral blood of piglets infected with porcine circovirus type 2. [Arch Virol. 163 \(5\): 1231-9.](#)

15. Liu, S. *et al.* (2019) Endothelial IL-8 induced by porcine circovirus type 2 affects dendritic cell maturation and antigen-presenting function. [Virol J. 16 \(1\): 154.](#)

16. Radlowski, E.C. *et al.* (2021) Combination-Feeding Causes Differences in Aspects of Systemic and Mucosal Immune Cell Phenotypes and Functions Compared to Exclusive Sow-Rearing or Formula-Feeding in Piglets. [Nutrients. 13\(4\):1097.](#)

17. Franzoni, G. *et al.* (2022) Analyses of the Impact of Immunosuppressive Cytokines on Porcine Macrophage Responses and Susceptibility to Infection to African Swine Fever Viruses. [Pathogens. 11 \(2\): 166.](#)

---

#### Further Reading

1. Piriou-Guzylack, L. (2008) Membrane markers of the immune cells in swine: an update. [Vet Res. 39: 54.](#)

---

#### Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in

frost-free freezers is not recommended.

<b>Guarantee</b>	12 months from date of despatch
<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10040 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA2314GA">https://www.bio-rad-antibodies.com/SDS/MCA2314GA</a> 10040
<b>Regulatory</b>	For research purposes only

## Related Products

### Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)	<a href="#">RPE</a>
Goat Anti Mouse IgG IgA IgM (STAR87...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (STAR76...)	<a href="#">RPE</a>
Goat Anti Mouse IgG (STAR70...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (H/L) (STAR117...)	<a href="#">Alk. Phos.</a> , <a href="#">DyLight®488</a> , <a href="#">DyLight®550</a> , <a href="#">DyLight®650</a> , <a href="#">DyLight®680</a> , <a href="#">DyLight®800</a> , <a href="#">FITC</a> , <a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR9...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (STAR77...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (Fc) (STAR120...)	<a href="#">FITC</a> , <a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR13...)	<a href="#">HRP</a>

### Recommended Negative Controls

[MOUSE IgG2b NEGATIVE CONTROL \(MCA691\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------	------------------	-------------------------------------------------------------------------------------------------------------------------------------------------	---------------	-----------------------------------------------------------------------------------------------------------------------------------------------------

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)  
'M383680:210513'

Printed on 13 Mar 2024