

Datasheet: MCA2317F

BATCH NUMBER 164573

Description:	MOUSE ANTI PIG MACROPHAGES:FITC
Specificity:	MACROPHAGES
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	BA4D5
Isotype:	IgG2b
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry (1)	▪			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) Membrane permeabilization is required for this application. The use of Leucoperm (Product Code [BUF09](#)) is recommended for this purpose.

Target Species	Pig		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant.		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% sodium azide (NaN ₃)		
Stabilisers	1% bovine serum albumin		

Approx. Protein Concentrations	IgG concentration 0.1 mg/ml
Immunogen	Porcine alveolar macrophages.
RRID	AB_567110
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the mouse SP2/0 mouse myeloma cell line.
Specificity	<p>Mouse anti Pig Macrophages antibody, clone BA4D5 recognizes porcine cells of the monocyte/macrophage lineage. Expression of the antigen is increased with maturation, with higher expression on peritoneal and alveolar macrophages.</p> <p>Some expression has also been observed on peripheral blood lymphocytes.</p> <p>The antigen recognized by clone BA4D5 has a broad tissue distribution and this antibody stains macrophages in a range of tissues, including the thymus, spleen periarteriolar lymphoid sheath (PALS), spleen red pulp and the Peyer's patches. Expression has also been reported on some non-hematopoietic cells including endothelial cells.</p> <p>It is believed that clone BA4D5 may be specific for porcine CD68 (Poulsen <i>et al.</i> 2016) although the protein recognized by this antibody has not yet been fully characterized. The protein is expressed on the cell surface, although it is most abundantly expressed in the cytoplasm.</p>
Flow Cytometry	Use 10µl of the suggested working dilution to 1x10 ⁶ cells in 100µl
References	<ol style="list-style-type: none"> Luechtenborg, B. <i>et al.</i> (2008) Function of scavenger receptor class A type I/II is not important for smooth muscle foam cell formation. Eur J Cell Biol. 87: 91-9. Ezquerro, A. <i>et al.</i> (2009) Porcine myelomonocytic markers and cell populations. Dev Comp Immunol. 33 (3): 284-98. Muscari C <i>et al.</i> (2010) Comparison between Culture Conditions Improving Growth and Differentiation of Blood and Bone Marrow Cells Committed to the Endothelial Cell Lineage. Biol Proced Online. 12 (1): 9023. Fujita M <i>et al.</i> (2013) Technique of endoscopic biopsy of islet allografts transplanted into the gastric submucosal space in pigs. Cell Transplant. 22 (12): 2335-44. Sohn, E.H. <i>et al.</i> (2015) Allogenic iPSC-derived RPE cell transplants induce immune response in pigs: a pilot study. Sci Rep. 5: 11791. Liu, G. <i>et al.</i> (2015) Influenza A Virus Panhandle Structure is Directly Involved in RIG-I Activation and IFN Induction. J Virol. pii: JVI.00232-15. Poulsen, C.B. <i>et al.</i> (2016) Treatment with a human recombinant monoclonal IgG antibody against oxidized LDL in atherosclerosis-prone pigs reduces cathepsin S in coronary lesions. Int J Cardiol. 215: 506-515. Rayat, G.R. <i>et al.</i> (2016) First update of the International Xenotransplantation Association consensus statement on conditions for undertaking clinical trials of porcine islet products in type 1 diabetes - Chapter 3: Porcine islet product manufacturing and release testing criteria. Xenotransplantation. 23 (1): 38-45.

9. Wang, L. *et al.* (2017) Porcine alveolar macrophage polarization is involved in inhibition of porcine reproductive and respiratory syndrome virus (PRRSV) replication. [J Vet Med Sci. 79 \(11\): 1906-15.](#)
10. Porras, A.M. *et al.* (2018) Creation of disease-inspired biomaterial environments to mimic pathological events in early calcific aortic valve disease. [Proc Natl Acad Sci U S A. 115 \(3\): E363-E371.](#)
11. Maciag, S.S. *et al.* (2022) On the influence of the source of porcine colostrum in the development of early immune ontogeny in piglets. [Sci Rep. 12 \(1\): 15630.](#)
12. dos Santos, M.C. *et al.* (2023) Effect of yeast extracted β -glucans on the immune response and reproductive performance of gilts in the adaptation, gestation, and lactation periods [Livestock Science. 275: 105289.](#)
13. Haach, V. *et al.* (2023) A polyvalent virosomal influenza vaccine induces broad cellular and humoral immunity in pigs. [Virology. 20 \(1\): 181.](#)
14. Petitpas, K. *et al.* (2022) Genetic modifications designed for xenotransplantation attenuate sialoadhesin-dependent binding of human erythrocytes to porcine macrophages. [Xenotransplantation. 29 \(6\): e12780.](#)
15. Forner, R. *et al.* (2021) Distribution difference of colostrum-derived B and T cells subsets in gilts and sows. [PLoS One. 16 \(5\): e0249366.](#)

Further Reading 1. Piriou-Guzylack, L. (2008) Membrane markers of the immune cells in swine: an update. [Vet Res. 39: 54.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10041 available at: <https://www.bio-rad-antibodies.com/SDS/MCA2317F>

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG2b NEGATIVE CONTROL:FITC \(MCA691F\)](#)

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M414441:221207'

Printed on 10 May 2026
