

Datasheet: MCA2317F

| | |
|----------------------|---------------------------------|
| Description: | MOUSE ANTI PIG MACROPHAGES:FITC |
| Specificity: | MACROPHAGES |
| Format: | FITC |
| Product Type: | Monoclonal Antibody |
| Clone: | BA4D5 |
| Isotype: | IgG2b |
| Quantity: | 0.1 mg |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|--------------------|-----|----|----------------|--------------------|
| Flow Cytometry (1) | ▪ | | | Neat |

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1) Membrane permeabilisation is required for this application. Bio-Rad recommends the use of Leucoperm™ (Product Code [BUF09](#)) for this purpose.

| | | | |
|---------------------------------------|---|----------------------------|--------------------------|
| Target Species | Pig | | |
| Product Form | Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid | | |
| Max Ex/Em | Fluorophore | Excitation Max (nm) | Emission Max (nm) |
| | FITC | 490 | 525 |
| Preparation | Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant | | |
| Buffer Solution | Phosphate buffered saline | | |
| Preservative Stabilisers | 0.09% Sodium Azide 1% Bovine Serum Albumin | | |
| Approx. Protein Concentrations | IgG concentration 0.1 mg/ml | | |

| | |
|------------------------|---|
| Immunogen | Porcine alveolar macrophages. |
| RRID | AB_567110 |
| Fusion Partners | Spleen cells from immunized BALB/c mice were fused with cells of the mouse SP2/0 mouse myeloma cell line. |
| Specificity | <p>Mouse anti Pig Macrophages antibody, clone BA4D5 recognizes porcine cells of the monocyte/macrophage lineage. Expression of the antigen is increased with maturation, with higher expression on peritoneal and alveolar macrophages.</p> <p>Some expression has also been observed on peripheral blood lymphocytes.</p> <p>The antigen recognized by clone BA4D5 has a broad tissue distribution and this antibody stains macrophages in a range of tissues, including the thymus, spleen periarteriolar lymphoid sheath (PALS), spleen red pulp and the Peyer's patches. Expression has also been reported on some non-hematopoietic cells including endothelial cells.</p> <p>It is believed that clone BA4D5 may be specific for porcine CD68 (Poulsen et al. 2016) although the protein recognized by this antibody has not yet been fully characterized. The protein is expressed on the cell surface, although it is most abundantly expressed in the cytoplasm.</p> |
| Flow Cytometry | Use 10ul of the suggested working dilution to 1x10 ⁶ cells in 100ul. |
| References | <ol style="list-style-type: none"> Luechtenborg, B. <i>et al.</i> (2008) Function of scavenger receptor class A type I/II is not important for smooth muscle foam cell formation. Eur J Cell Biol. 87: 91-9. Fujita M <i>et al.</i> (2013) Technique of endoscopic biopsy of islet allografts transplanted into the gastric submucosal space in pigs. Cell Transplant. 22 (12): 2335-44. Muscari C <i>et al.</i> (2010) Comparison between Culture Conditions Improving Growth and Differentiation of Blood and Bone Marrow Cells Committed to the Endothelial Cell Lineage. Biol Proced Online. 12 (1): 9023. Liu, G. <i>et al.</i> (2015) Influenza A Virus Panhandle Structure is Directly Involved in RIG-I Activation and IFN Induction. J Virol. pii: JVI.00232-15. Ezquerro, A. <i>et al.</i> (2009) Porcine myelomonocytic markers and cell populations. Dev Comp Immunol. 33 (3): 284-98. Rayat, G.R. <i>et al.</i> (2016) First update of the International Xenotransplantation Association consensus statement on conditions for undertaking clinical trials of porcine islet products in type 1 diabetes - Chapter 3: Porcine islet product manufacturing and release testing criteria. Xenotransplantation. 23 (1): 38-45. Poulsen, C.B. <i>et al.</i> (2016) Treatment with a human recombinant monoclonal IgG antibody against oxidized LDL in atherosclerosis-prone pigs reduces cathepsin S in coronary lesions. Int J Cardiol. 215: 506-515. Sohn, E.H. <i>et al.</i> (2015) Allogenic iPSC-derived RPE cell transplants induce immune response in pigs: a pilot study. Sci Rep. 5: 11791. Wang, L. <i>et al.</i> (2017) Porcine alveolar macrophage polarization is involved in inhibition of porcine reproductive and respiratory syndrome virus (PRRSV) replication. J Vet Med Sci. Sep 17 [Epub ahead of print]. |

10. Porras, A.M. *et al.* (2018) Creation of disease-inspired biomaterial environments to mimic pathological events in early calcific aortic valve disease. [Proc Natl Acad Sci U S A. 115 \(3\): E363-E371.](#)

Further Reading 1. Piriou-Guzylack, L. (2008) Membrane markers of the immune cells in swine: an update. [Vet Res. 39: 54.](#)

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10041 available at: 10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG2b NEGATIVE CONTROL:FITC \(MCA691F\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

From March 15, 2021, we will no longer supply printed datasheets with our products.

Look out for updates on how to access your digital version at bio-rad-antibodies.com

'M366639:200529'

Printed on 12 Feb 2021

© 2021 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)