

Datasheet: MCA2318EL BATCH NUMBER 148129

Description:	MOUSE ANTI HUMAN CD209:Low Endotoxin
Specificity:	CD209
Other names:	DC-SIGN
Format:	Low Endotoxin
Product Type:	Monoclonal Antibody
Clone:	MR-1
Isotype:	lgG1
Quantity:	0.5 mg

Product Details

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u>.

		Yes	No	Not Determined	Suggested Dilution
	Flow Cytometry	•			
	Immunohistology - Frozen			•	
	Immunohistology - Paraffin			•	
	ELISA			•	
	Immunoprecipitation			•	
	Western Blotting			•	
	Immunofluorescence	-			
	Functional Assays	-			
	Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.				technique this does not
					•
Target Species	Human				
Product Form	Purified IgG - liquid				
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant				
Buffer Solution	Phosphate buffered saline				

Preservative Stabilisers	None present			
Carrier Free	Yes			
Endotoxin Level	< 0.01 EU/ug			
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml			
Immunogen	Immature myeloid monocyte-derived dendritic cells (MDDCs).			
External Database Links	UniProt: <u>Q9NNX6</u> <u>Related reagents</u> Entrez Gene: <u>30835</u> CD209 <u>Related reagents</u>			
Synonyms	CLEC4L			
RRID	AB_566636			
Fusion Partners	Spleen cells from immunized Balb/c mice were fuzed with cells of the Mouse SP2/0-Ag14 myeloma cell line.			
Specificity	Mouse anti Human CD209 antibody, clone MR-1 recognizes human DC-specific ICAM-3 grabbing nonintegrin (DC-SIGN), a ~45 kDa C-type lectin that binds ICAM-3 also known as CD209. CD209 is primarily expressed on a population of immature dendritic cells in peripheral tissues and on immature myeloid monocyte-derived DC's <i>in vitro</i> . CD209 is involved in dendritic cell (DC) migration and the initial interaction between DC's and naive T lymphocytes. CD209 also binds HIV-1 gp120 and plays a key role in the dissemination of HIV-1 by DC's. Mouse anti Human CD209 antibody, clone MR-1 is reported to partially block the functional activity of DC-SIGN (<u>Melero <i>et al.</i></u>).			
	cells in peripheral tissues and on immature myeloid monocyte-derived DC's <i>in vitro</i> . CD209 is involved in dendritic cell (DC) migration and the initial interaction between DC's and naive T lymphocytes. CD209 also binds HIV-1 gp120 and plays a key role in the dissemination of HIV-1 by DC's. Mouse anti Human CD209 antibody, clone MR-1 is reported to partially block the			
Flow Cytometry	cells in peripheral tissues and on immature myeloid monocyte-derived DC's <i>in vitro</i> . CD209 is involved in dendritic cell (DC) migration and the initial interaction between DC's and naive T lymphocytes. CD209 also binds HIV-1 gp120 and plays a key role in the dissemination of HIV-1 by DC's. Mouse anti Human CD209 antibody, clone MR-1 is reported to partially block the			

	 5. Alvarez, C.P. <i>et al</i> (2002) C-type lectins DC-SIGN and L-SIGN mediate cellular er Ebola virus in cis and in trans. J Virol. 76: 6841-4. 6. Zhang, S.S. <i>et al.</i> (2008) Plasminogen activator Pla of Yersinia pestis utilizes mur DEC-205 (CD205) as a receptor to promote dissemination J Biol Chem. 283: 31511 7. Relloso, M. <i>et al.</i> (2002) Dendritic cell (DC)-specific intercellular adhesion molecul (ICAM-3)-grabbing nonintegrin (DC-SIGN, CD209), a C-type surface lectin in humar is a receptor for Leishmania amastigotes. J Biol Chem. 277:36766-9 8. Ciudad, M.T. <i>et al.</i> (2017) Analysis of the HLA-DR peptidome from human dendritic cells reveals high affinity repertoires and nonconventional pathways of peptide general Leukoc Biol. 101 (1): 15-27. 9. Chen, J.M. <i>et al.</i> (2017) Bovine Lactoferrin Inhibits Dengue Virus Infectivity by Interacting with Heparan Sulfate, Low-Density Lipoprotein Receptor, and DC-SIGN. Mol Sci. 18 (9)Sep 12 [Epub ahead of print]. 	ine - <u>21.</u> le 3 n DCs, ic ration.
Storage	Store at -20°C only. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.	
Guarantee	12 months from date of despatch	
Health And Safety Information	Material Safety Datasheet documentation #10162 available at: https://www.bio-rad-antibodies.com/SDS/MCA2318EL 10162	
Regulatory	For research purposes only	

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12)	RPE			
Goat Anti Mouse IgG IgA IgM (STAR87) <u>HRP</u>				
Goat Anti Mouse IgG (STAR76)	RPE			
Rabbit Anti Mouse IgG (STAR13)	HRP			
Goat Anti Mouse IgG (STAR70)	<u>FITC</u>			
Goat Anti Mouse IgG (H/L) (STAR117)	Alk. Phos., DyLight®488, DyLight®550,			
	DyLight®650, DyLight®680, DyLight®800,			
	FITC, HRP			
Rabbit Anti Mouse IgG (STAR9)	<u>FITC</u>			
Goat Anti Mouse IgG (STAR77)	HRP			
Goat Anti Mouse IgG (Fc) (STAR120)	FITC, HRP			
Recommended Negative Controls				

MOUSE IgG1 NEGATIVE CONTROL:Low Endotoxin (MCA928EL)

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

North & South Tel: +1 800 265 7376

Worldwide

Tel: +44 (0)1865 852 700 Europe Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com

Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint

Printed on 19 Jan 2024

America

Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com 'M366642:200529'