

Datasheet: MCA2318PET

Description:	MOUSE ANTI HUMAN CD209:RPE
Specificity:	CD209
Other names:	DC-SIGN
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	MR-1
Isotype:	IgG1
Quantity:	25 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized		
Reconstitution	Reconstitute in 0.25 ml distilled water		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE 488nm laser	496	578
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide		
Stabilisers	1% Bovine Serum Albumin		
	5% Sucrose		
Immunogen	Immature myeloid monocyte-derived dendritic cells (MDDCs).		
External Database Links	UniProt:		
	Q9NNX6	Related reagents	

Entrez Gene:[30835](#) CD209 [Related reagents](#)

Synonyms	CLEC4L
-----------------	--------

Fusion Partners	Spleen cells from immunized Balb/c mice were fused with cells of the Mouse SP2/0-Ag14 myeloma cell line.
------------------------	--

Specificity	<p>Mouse anti Human CD209 antibody, clone MR-1 recognizes human DC-specific ICAM-3 grabbing nonintegrin (DC-SIGN), a ~45 kDa C-type lectin that binds ICAM-3 also known as CD209. CD209 is primarily expressed on a population of immature dendritic cells in peripheral tissues and on immature myeloid monocyte-derived DC's <i>in vitro</i>.</p> <p>CD209 is involved in dendritic cell (DC) migration and the initial interaction between DC's and naive T lymphocytes. CD209 also binds HIV-1 gp120 and plays a key role in the dissemination of HIV-1 by DC's.</p> <p>Mouse anti Human CD209 antibody, clone MR-1 is reported to partially block the functional activity of DC-SIGN (Melero et al.).</p>
--------------------	---

Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
-----------------------	---

References	<ol style="list-style-type: none">1. Relloso, M. <i>et al.</i> (2002) DC-SIGN (CD209) expression is IL-4 dependent and is negatively regulated by IFN, TGF-beta, and anti-inflammatory agents. J Immunol. 168 (6): 2634-43.2. Melero, I. <i>et al.</i> (2002) An anti-ICAM-2 (CD102) monoclonal antibody induces immune-mediated regressions of transplanted ICAM-2-negative colon carcinomas. Cancer Res. 62 (11): 3167-74.3. Domínguez-Soto, A. <i>et al.</i> (2011) Dendritic Cell-Specific ICAM-3-Grabbing Nonintegrin Expression on M2-Polarized and Tumor-Associated Macrophages Is Macrophage-CSF Dependent and Enhanced by Tumor-Derived IL-6 and IL-10. J Immunol. 186: 2192-200.4. Chang, S.K. <i>et al.</i> (2008) B lymphocyte stimulator regulates adaptive immune responses by directly promoting dendritic cell maturation. J Immunol. 180: 7394-403.5. Alvarez, C.P. <i>et al.</i> (2002) C-type lectins DC-SIGN and L-SIGN mediate cellular entry by Ebola virus in cis and in trans. J Virol. 76: 6841-4.6. Zhang, S.S. <i>et al.</i> (2008) Plasminogen activator Pla of <i>Yersinia pestis</i> utilizes murine DEC-205 (CD205) as a receptor to promote dissemination J Biol Chem. 283: 31511-21.7. Relloso, M. <i>et al.</i> (2002) Dendritic cell (DC)-specific intercellular adhesion molecule 3 (ICAM-3)-grabbing nonintegrin (DC-SIGN, CD209), a C-type surface lectin in human DCs, is a receptor for <i>Leishmania amastigotes</i>. J Biol Chem. 277:36766-98. Ciudad, M.T. <i>et al.</i> (2016) Analysis of the HLA-DR peptidome from human dendritic cells reveals high affinity repertoires and nonconventional pathways of peptide generation. J Leukoc Biol. Jun 30. pii: jlb.6HI0216-069R. [Epub ahead of print]9. Chen, J.M. <i>et al.</i> (2017) Bovine Lactoferrin Inhibits Dengue Virus Infectivity by Interacting with Heparan Sulfate, Low-Density Lipoprotein Receptor, and DC-SIGN. Int J Mol Sci. 18 (9)Sep 12 [Epub ahead of print].
-------------------	---

Storage	Store at +4°C.
	DO NOT FREEZE
	This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life	12 months from date of reconstitution.
-------------------	--

Health And Safety Information Material Safety Datasheet documentation #10075 available at:
10075: <https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf>

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:RPE \(MCA928PE\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M323656:180727'

Printed on 06 Dec 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)