

Datasheet: MCA2318T

BATCH NUMBER 149091

Description:	MOUSE ANTI HUMAN CD209
Specificity:	CD209
Other names:	DC-SIGN
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	MR-1
Isotype:	IgG1
Quantity:	25 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative	0.09% Sodium Azide

Stabilisers

Carrier Free Yes

Approx. Protein Concentrations IgG concentration 1.0 mg/ml

Immunogen Immature myeloid monocyte-derived dendritic cells (MDDCs).

External Database Links

UniProt:

[Q9NNX6](#) [Related reagents](#)

Entrez Gene:

[30835](#) CD209 [Related reagents](#)

Synonyms CLEC4L

RRID AB_1101360

Fusion Partners Spleen cells from immunized Balb/c mice were fused with cells of the Mouse SP2/0-Ag14 myeloma cell line.

Specificity **Mouse anti Human CD209 antibody, clone MR-1** recognizes human DC-specific ICAM-3 grabbing nonintegrin (DC-SIGN), a ~45 kDa C-type lectin that binds ICAM-3 also known as CD209. CD209 is primarily expressed on a population of immature dendritic cells in peripheral tissues and on immature myeloid monocyte-derived DC's *in vitro*. CD209 is involved in dendritic cell (DC) migration and the initial interaction between DC's and naive T lymphocytes. CD209 also binds HIV-1 gp120 and plays a key role in the dissemination of HIV-1 by DC's. Mouse anti Human CD209 antibody, clone MR-1 is reported to partially block the functional activity of DC-SIGN ([Melero *et al.*](#)).

Flow Cytometry Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

References

1. Relloso, M. *et al.* (2002) DC-SIGN (CD209) expression is IL-4 dependent and is negatively regulated by IFN, TGF-beta, and anti-inflammatory agents. [J Immunol. 168 \(6\): 2634-43.](#)
2. Melero, I. *et al.* (2002) An anti-ICAM-2 (CD102) monoclonal antibody induces immune-mediated regressions of transplanted ICAM-2-negative colon carcinomas. [Cancer Res. 62 \(11\): 3167-74.](#)
3. Domínguez-Soto, A. *et al.* (2011) Dendritic Cell-Specific ICAM-3-Grabbing Nonintegrin Expression on M2-Polarized and Tumor-Associated Macrophages Is Macrophage-CSF Dependent and Enhanced by Tumor-Derived IL-6 and IL-10. [J Immunol. 186: 2192-200.](#)
4. Chang, S.K. *et al.* (2008) B lymphocyte stimulator regulates adaptive immune responses by directly promoting dendritic cell maturation. [J Immunol. 180: 7394-403.](#)
5. Alvarez, C.P. *et al.* (2002) C-type lectins DC-SIGN and L-SIGN mediate cellular entry by Ebola virus in cis and in trans. [J Virol. 76: 6841-4.](#)
6. Zhang, S.S. *et al.* (2008) Plasminogen activator Pla of *Yersinia pestis* utilizes murine

- DEC-205 (CD205) as a receptor to promote dissemination [J Biol Chem. 283: 31511-21.](#)
7. Relloso, M. *et al.* (2002) Dendritic cell (DC)-specific intercellular adhesion molecule 3 (ICAM-3)-grabbing nonintegrin (DC-SIGN, CD209), a C-type surface lectin in human DCs, is a receptor for Leishmania amastigotes. [J Biol Chem. 277:36766-9](#)
8. Ciudad, M.T. *et al.* (2017) Analysis of the HLA-DR peptidome from human dendritic cells reveals high affinity repertoires and nonconventional pathways of peptide generation. [J Leukoc Biol. 101 \(1\): 15-27.](#)
9. Chen, J.M. *et al.* (2017) Bovine Lactoferrin Inhibits Dengue Virus Infectivity by Interacting with Heparan Sulfate, Low-Density Lipoprotein Receptor, and DC-SIGN. [Int J Mol Sci. 18 \(9\)Sep 12 \[Epub ahead of print\].](#)

Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA2318T 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M366645:200529'

