Datasheet: MCA2320

Description:	MOUSE ANTI HUMAN MAdCAM-1
Specificity:	MAdCAM-1
Other names:	MUCOSAL ADDRESSIN CELL ADHESION MOLECULE-1
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	314G8
Isotype:	IgG1
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Frozen	▪			1/25 - 1/200
Immunohistology - Paraffin			▪	
ELISA	▪			
Immunoprecipitation			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% sodium azide (NaN_3)
Carrier Free	Yes

Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Recombinant soluble MAdCAM-1-fc fusion protein.
External Database Links	**UniProt:** Q13477 Related reagents **Entrez Gene:** 8174 MADCAM1 Related reagents
RRID	AB_567120
Fusion Partners	Spleen cells from immunised Balb/c mice were fused with cells of the mouse P3X63Ag8.653 myeloma cell line.
Specificity	**Mouse anti Human MAdCAM-1 antibody, clone 314G8** recognizes human mucosal addressin cell adhesion molecule 1 (MAdCAM-1) a 60kD cell surface protein that is involved in lymphocyte trafficking. MAdCAM-1 is expressed on high endothelial venules of Peyer's patches and mesenteric lymph nodes. MAdCAM-1 expression has also been reported in the gut lamina propria but clone 314G8 does not recognize MAdCAM-1 in these tissues. Clone 314G8 reacts with the ligand-binding first Ig domain and reports suggest that a splice variant exists in the gut which is not recognized by clone 314G8. Clone 314G8 is reported to block the interaction of MAdCAM-1 with alpha 4 beta 7 (Leung *et al.* 2004).
Flow Cytometry	Use 10µl of the suggested working dilution to label 10^6 cells in 100µl
References	1. Leung, E. *et al.* (1996) Cloning of the mucosal addressin MAdCAM-1 from human brain: identification of novel alternatively spliced transcripts. Immunol Cell Biol. 74 (6): 490-6. 2. Leung, E. *et al.* (2004) Bioassay detects soluble MAdCAM-1 in body fluids. Immunol Cell Biol. 82 (4): 400-9. 3. Olloquequi, J. *et al.* (2011) Significant increase of CD57+ cells in pulmonary lymphoid follicles of COPD patients. Eur Respir J. 37 (2): 289-98. 4. Guerra-Pérez, N. *et al.* (2015) Retinoic Acid Imprints a Mucosal-like Phenotype on Dendritic Cells with an Increased Ability To Fuel HIV-1 Infection. J Immunol. pii: 1402623. 5. Goode, D. *et al.* (2014) Sex hormones selectively impact the endocervical mucosal microenvironment: implications for HIV transmission. PLoS One. 9 (5): e97767. 6. Guzzo, C. *et al.* (2017) Virion incorporation of integrin αα4β7 facilitates HIV-1 infection and intestinal homing. Sci Immunol. 2 (11) [Epub ahead of print].
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA2320 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos., HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos., DyLight®488, DyLight®550, DyLight®650, DyLight®680, DyLight®800, FITC, HRP
Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC, HRP

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	**Worldwide**	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	**Europe**	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M414508:221209'

Printed on 19 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint