

Datasheet: MCA2320EL

Description:	MOUSE ANTI HUMAN MAdCAM-1:Low Endotoxin		
Specificity:	MAdCAM-1		
Other names:	MUCOSAL ADDRESSIN CELL ADHESION MOLECULE-1		
Format:	Low Endotoxin		
Product Type:	Monoclonal Antibody		
Clone:	314G8		
Isotype:	lgG1		
Quantity:	0.5 mg		

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal					
	communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-</u>					
	rad-antibodies.com/protocols.					
	Tad-antibodies.com/proto	Yes No Not Determined				
	Immunohistology - Frozen			•	Suggested Dilution	
	Immunohistology - Paraffin					
	ELISA					
	Immunoprecipitation			•		
	Functional Assays	-				
	Where this product has n	ot been t	ested for	use in a particular tech	inique this does not	
	necessarily exclude its use in such procedures. Suggested working dilutions are given a a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.					
Target Species	Human					
Product Form	Purified IgG - liquid					
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant					
Buffer Solution	Phosphate buffered saline					
Preservative Stabilisers	None present					
Carrier Free	Yes					

Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Recombinant soluble MAdCAM-1-fc fusion protein.
External Database Links	UniProt: <u>Q13477</u> <u>Related reagents</u> Entrez Gene: <u>8174</u> MADCAM1 <u>Related reagents</u>
RRID	AB_567122
Fusion Partners	Spleen cells from immunised Balb/c mice were fused with cells of the mouse P3X63Ag8.653 myeloma cell line.
Specificity	Mouse anti Human MAdCAM-1 antibody, clone 314G8 recognizes human mucosal addressin cell adhesion molecule 1 (MAdCAM-1) a 60kD cell surface protein that is involved in lymphocyte trafficking. MAdCAM-1 is expressed on high endothelial venules of Peyer's patches and mesenteric lymph nodes. MAdCAM-1 expression has also been reported in the gut lamina propria but clone 314G8 does not recognize MAdCAM-1 in these tissues. Clone 314G8 reacts with the ligand-binding first Ig domain and reports suggest that a splice variant exists in the gut which is not recognized by clone 314G8. Clone 314G8 is reported to block the interaction of MAdCAM-1 with alpha 4 beta 7 (Leung et al. 2004).
Flow Cytometry	Use 10μ I of the suggested working dilution to label 10^6 cells in 100μ I
References	 Leung, E. <i>et al.</i> (1996) Cloning of the mucosal addressin MAdCAM-1 from human brain: identification of novel alternatively spliced transcripts. <u>Immunol Cell Biol. 74 (6): 490-6.</u> Leung, E. <i>et al.</i> (2004) Bioassay detects soluble MAdCAM-1 in body fluids. <u>Immunol Cell Biol. 82 (4): 400-9.</u> Olloquequi, J. <i>et al.</i> (2011) Significant increase of CD57+ cells in pulmonary lymphoid follicles of COPD patients. <u>Eur Respir J. 37 (2): 289-98.</u> Guerra-Pérez, N. <i>et al.</i> (2015) Retinoic Acid Imprints a Mucosal-like Phenotype on Dendritic Cells with an Increased Ability To Fuel HIV-1 Infection. <u>J Immunol. pii: 1402623.</u> Goode, D. <i>et al.</i> (2014) Sex hormones selectively impact the endocervical mucosal microenvironment: implications for HIV transmission. <u>PLoS One. 9 (5): e97767.</u> Guzzo, C. <i>et al.</i> (2017) Virion incorporation of integrin αα4β7 facilitates HIV-1 infection and intestinal homing. <u>Sci Immunol. 2 (11) [Epub ahead of print].</u>
Storage	Store at -20°C. This product should be stored undiluted. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10162 available at: https://www.bio-rad-antibodies.com/SDS/MCA2320EL 10162
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77)	HRP			
Rabbit Anti Mouse IgG (STAR12)	RPE			
Goat Anti Mouse IgG (STAR70)	<u>FITC</u>			
Goat Anti Mouse IgG IgA IgM (STAR87) <u>Alk. Phos.</u> , <u>HRP</u>				
Rabbit Anti Mouse IgG (STAR9)	<u>FITC</u>			
Goat Anti Mouse IgG (STAR76)	RPE			
Goat Anti Mouse IgG (H/L) (STAR117)	Alk. Phos., DyLight®488, DyLight®550,			
	DyLight®650, DyLight®680, DyLight®800,			
	FITC, HRP			
Rabbit Anti Mouse IgG (STAR13)	HRP			
Goat Anti Mouse IgG (Fc) (STAR120)	FITC, HRP			
Recommended Negative Controls				

MOUSE IgG1 NEGATIVE CONTROL:Low Endotoxin (MCA928EL)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-ra	ad.com	Email: antibody_sales_uk@bio-r	ad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M414507:221209'

Printed on 25 Oct 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint