

Datasheet: MCA2332T

Description:	MOUSE ANTI HUMAN CD261		
Specificity:	CD261		
Other names:	DR4, TRAIL-R1		
Format:	Purified		
Product Type:	Monoclonal Antibody		
Clone:	DR-4-02		
Isotype:	lgG1		
Quantity:	25 μg		

Product Details

RRID AB_1101654

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			1/50 - 1/100
Immunohistology - Frozen			•	
Immunohistology - Paraffin			•	
ELISA			•	
Immunoprecipitation	-			
Western Blotting			•	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human	
Product Form	Purified IgG - liquid	
Preparation	Purified IgG prepared by affinity chromatography on Protein A from	tissue culture supernatant
Buffer Solution	Phosphate buffered saline	
Preservative Stabilisers	0.09% Sodium Azide	
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml	
Immunogen	Fusion protein containing the extracellular region of CD261 (DR4).	

External Database Links	UniProt:			
	O00220 Related reagents			
	Entrez Gene:			
	8797 TNFRSF10A Related reagents			
Synonyms	APO2, DR4, TRAILR1			
Fusion Partners	Spleen cells from immunized F1 hybrid mice were fuzed with cells of the SP2/0-Ag14 myeloma cell line.			
Specificity	Mouse anti Human CD261 antibody, clone DR-4-02 recognizes human death receptor 4 (DR4), also known as CD261 or TRAIL- R1. DR4 is a type I transmembrane protein of 468 amino acids, which is expressed in most human tissues including spleen, peripheral blood leucocytes, thymus and in a variety of tumour-derived cell lines.			
	DR4 plays a role in inducing cell death. The binding of TRAIL to DR4 triggers the activation of pro-caspases 8 and 10, leading to apoptosis.			
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.			
References	1. Cosgrove, C. <i>et al.</i> (2013) Early and nonreversible decrease of CD161++/MAIT cells in HIV infection. <u>Blood. 121: 951-61.</u>			
	2. Crescenzi, E. et al. (2011) NF-κB-dependent cytokine secretion controls Fas expression on			
	chemotherapy-induced premature senescent tumor cells. Oncogene. 30: 2707-17.			
	3. Zhang, Y. and Zhang, B. (2008) TRAIL resistance of breast cancer cells is associated with			
	constitutive endocytosis of death receptors 4 and 5. Mol Cancer Res. 6: 1861-71.			
Storage	Store at +4°C or at -20°C if preferred.			
	This product should be stored undiluted.			
	Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.			

Regulatory

Health And Safety

Shelf Life

Information

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG IgA IgM (STAR87...) Alk. Phos., HRP

18 months from date of despatch.

For research purposes only

Material Safety Datasheet documentation #10040 available at:

10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf

Goat Anti Mouse IgG (STAR77...) <u>HRP</u>

Rabbit Anti Mouse IgG (STAR12...) RPE

Rabbit Anti Mouse IgG (STAR8...) <u>DyLight®800</u>

Rabbit Anti Mouse IgG (STAR13...)

Goat Anti Mouse IgG (STAR76...)

Goat Anti Mouse IgG (STAR70...)

FITC

Goat Anti Mouse IgG (Fc) (STAR120...)

FITC, HRP

Rabbit Anti Mouse IgG (STAR9...) FITC

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®549,

<u>DyLight®649</u>, <u>DyLight®680</u>, <u>DyLight®800</u>,

FITC, HRP

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

North & South Tel: +1 800 265 7376 Worldwide

Fax: +1 919 878 3751

America

Email: antibody_sales_us@bio-rad.com

Tel: +44 (0)1865 852 700 **Europe** Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M346237:190201'

Printed on 01 Feb 2019

© 2019 Bio-Rad Laboratories Inc | Legal | Imprint